

Avifaunal surveys in the central Peruvian Amazon clarify range limits and highlight links between avian and habitat diversity

Andre E. Moncrieff,^{1*} Oscar Johnson,¹ Cristhian Felix,² Anna E. Hiller,¹ Eamon C. Corbett,¹ Matthew L. Brady,¹ Glenn F. Seeholzer,^{1,3} Emil Bautista,² Daniel F. Lane,¹ and Michael G. Harvey^{1,4}

ABSTRACT—The bird life of the central Peruvian Amazon is poorly known. To better characterize bird diversity and distributions, we conducted 4 expeditions to this lowland area, totaling 114 field days in 2015, 2018, and 2019. We focused on sampling under-surveyed habitats, *terra firme* in multiple interfluvies, and sites around a recent river avulsion, and we detected 657 bird species across 22 study sites. Our results include the first extensive inventories of blackwater wetland systems, *Guadua* bamboo tracts, and riverine islands in central Peru; revised information on distributions with respect to river barriers, including documentation of contact and putative hybridization between parapatric forms; 28 first regional records; and new details on poorly known taxa. Together, these results provide a more complete picture of avian diversity in an area characterized by high species richness, high habitat diversity, and biogeographic interchange. Received 30 June 2020. Accepted 4 March 2021.

Key words: Amazonia, biodiversity, biogeography, contact zone, hybridization, river avulsion, Ucayali.

Reconocimientos de avifauna de la Amazonía central del Perú clarifican límites en distribuciones y destacan el vínculo entre diversidad aviar y de hábitats

RESUMEN (Spanish)—La avifauna de la Amazonía central del Perú es pobremente conocida. Para caracterizar su diversidad y distribución, llevamos a cabo 4 expediciones en esta área de tierras bajas, con un total de 114 días de trabajo de campo en 2015, 2018 y 2019. Nos enfocamos en muestrear hábitats subestudiados, *terra firme* en múltiples interfluvios y sitios alrededor de avulsiones recientes de ríos. Detectamos 657 especies de aves en 22 sitios de estudio. Nuestros resultados incluyen los primeros inventarios extensos de sistemas de humedales de aguas negras, parches de bambú *Guadua* e islas en los ríos del Perú central; información corregida de distribuciones respecto a barreras ribereñas, incluida documentación de contacto e hibridación putativa entre formas parapatricas; 28 primeros registros regionales y nuevos detalles de taxa escasamente conocidos. En conjunto, estos resultados proveen una imagen más completa de la diversidad aviar en un área caracterizada por una alta riqueza de especies, una alta diversidad de hábitats e intercambio biogeográfico.

Palabras clave: Amazonia, avulsión de ríos, biodiversidad, biogeografía, hibridación, Ucayali, zona de contacto.

The central Peruvian Amazon lies within a global hotspot of bird species richness in the western Amazon Basin (Orme et al. 2005). This richness is tied to the diversity of avian habitats regionally (Terborgh 1985, Parker et al. 1996), including upland forest (*terra firme*), floodplain forest along silt-laden rivers draining out of the Andes (*várzea*), blackwater drainages with associated seasonally flooded forests (*igapó*), wetlands, riverine habitats, and associated islands (Remsen and Parker 1983, Rosenberg 1990), *Guadua* (Poaceae) bamboo tracts (Kratte 1997, Parker et al. 1997, Lebbin 2007), stunted forest on nutrient-poor soils (Álvarez et al. 2013, Borges et al. 2016),

and, increasingly, agricultural land and settlements. Understanding how spatial patterns of avian diversity correspond to habitat diversity helps to clarify the ecological and evolutionary basis of diversity hotspots (Remsen 1985, Terborgh 1985, Marra and Remsen 1997). However, more information is needed on avian habitat associations and on the geographic distributions of habitat specialists in the western Amazon Basin.

The high species richness of the western Amazon Basin is also linked to avian biogeographic turnover across major landscape barriers such as rivers. The central Peruvian Amazon is at the headwaters of the Amazon River and at the confluence of avifaunas from the Napo area of endemism north of the Amazon and the Inambari area of endemism to the south (Cracraft 1985, Silva et al. 2005). Identifying the locations of range limits and the presence or absence of contact between closely related taxa in headwater regions can clarify not only patterns of species richness, but also how Amazonian species form and are maintained (Bates et al. 2004, Naka et al. 2012,

¹ Department of Biological Sciences and Museum of Natural Science, Louisiana State University, Baton Rouge, LA, USA

² Centro de Ornitología y Biodiversidad, Urb. Huertos de San Antonio, Surco, Lima, Perú

³ Department of Ornithology, American Museum of Natural History, New York, NY, USA

⁴ Department of Biological Sciences and Biodiversity Collections, The University of Texas at El Paso, TX, USA

* Corresponding author: amoncr2@lsu.edu

Weir et al. 2015, Naka and Brumfield 2018). The central Peruvian Amazon is biogeographically complex, with the Ucayali, Urubamba, and Tambo rivers as well as outlying mountain ranges like the Cerros del Sira serving as distributional barriers. These barriers, moreover, are not fixed over time. The Ucayali River in particular is thought to have had a dynamic history even over recent timescales (Lathrap 1968, Dumont 1991), with a notable example being an avulsion that passively transferred ~2,000 km² of land near Pucallpa from the west to the east side of the river in the late 18th century (hereafter, the Ucayali avulsion; Pärssinen et al. 1996). High-resolution survey data are required to assess the role of these landscape barriers in shaping geographic distributions and contact zones between taxa.

The first major ornithological surveys of the central Peruvian Amazon were conducted in 1927–1928 by the Olalla brothers, Alfonso and Ramón, who collected ~2,000 specimens along the upper Ucayali River (Wiley 2010). Further surveys of similar scale by the Louisiana State University Museum of Natural Science (LSUMNS), Museo de Historia Natural de la Universidad Nacional Mayor de San Marcos (UNMSM), and Centro de Ornitología y Biodiversidad (CORBIDI) have established general biogeographic patterns. These surveys include those at Yarinacocha from 1961 to 1972 (~900 specimens; O'Neill and Pearson 1974), near Balta in the upper Purús River drainage from 1963 to 1971 (~2,300 specimens; Lowery and O'Neill 1965, 1966, 1969; O'Neill 1969, 1974, 2003), along the Shesha River in 1987 in what is now the Sierra del Divisor National Park (park established in 2015; ~1,100 specimens; O'Neill et al. 1991; also see Schulenberg et al. 2006), and to sites around the upper Ucayali River in 2010–2011, particularly along the Cohengua River (~2,100 specimens; Harvey et al. 2014b). Additional surveys around Pucallpa (Traylor 1958, Dickerman 1975, Srinivas and Molina 2013, Srinivas and Koh 2016), the upper Purús River drainage (Angulo et al. 2016), upper Jurua drainage (Balta 2007), Mapuya and Caco Rivers (Goussard 1984), and lower Urubamba River (M.J. Miller, University of Oklahoma, unpubl. data), and more peripherally in the foothills of the Cerros del Sira (Terborgh and Weske 1975, González 1998, Mee et al. 2002, Harvey et al. 2011, Socolar et al. 2013), Gran Pajonal (Harvey et al. 2011), northern

foothills of the Cordillera Vilcabamba (Schulenberg and Servat 2001), and lowlands of Acre, Brazil (Whittaker and Oren 1999), provided important context for the work presented here.

Despite this long history, ornithological study in the central Peruvian Amazon has been patchy both in time and space. The area generally has not been the focus of intensive avifaunal surveys such as those conducted in the Loreto and Madre de Dios regions by ornithologists and amateurs alike (e.g., Graham et al. 1980, Parker 1982, Capparella 1987, Rosenberg 1990, Terborgh et al. 1990, Robinson 1997, Lebbin 2007, Álvarez et al. 2013, Díaz-Alván et al. 2017, Schmitt et al. 2017, Williams 2017, Socolar et al. 2018, Moncrieff et al. 2019), certainly in part due to much less developed infrastructure such as roads, research centers, and eco-lodges. Major habitats, particularly extensive blackwater wetlands and surrounding *igapó* forests, *Guadua* bamboo tracts, and riverine islands, remain largely unexplored. Knowledge of bird distributions with respect to rivers and the extent of hybridization between closely related taxa are also poorly known (but see Harvey et al. 2014b). Meanwhile, expanding human settlement is providing increased access, but threatens the very habitats in need of study.

Here, we report on surveys aimed at filling these avian knowledge gaps in the central Peruvian Amazon. We conducted the most extensive surveys to date of blackwater wetland systems, *Guadua* bamboo tracts, and riverine islands in the area, and targeted specific localities to assess the influence of the Ucayali River and its tributaries on the geographic ranges of birds, including sites within the ~2,000 km² land mass that switched riverbanks during the Ucayali avulsion.

Methods

Study sites

We surveyed 22 sites across central Peru over 114 d during 2015, 2018, and 2019 (Table 1; Fig. 1). The study area is largely located within the Ucayali Region, but also includes lowland sites in the Huánuco, Junín, and Cuzco regions (we restrict our usage of “region” to the first-level administrative subdivision in Peru previously known as a “department”). Sampling effort was focused along the Ucayali River and its main tributaries, the Urubamba and Tambo rivers, which join to form

Table 1. Summary of avifaunal surveys in the central Amazon of Peru.

Study sites	Dates surveyed	Surveyors	Species detected	Specimens collected	Species collected	Species with media ^a
Yarinacocha	19 Dec 2015; 25, 27 Jul 2019; 20, 31 Aug–1 Sep 2019	AEM, OJ, CFC, ECC, MLB, MGH	112	2	1	45
Pucallpa area	12 Dec 2015; 24–26 Jul 2019; 4, 7, 25, 29 Aug 2019; 1 Sep 2019	AEM, OJ, CFC, ECC, MLB, MGH, DFL	92	2	1	5
Tournavista area	2, 25–29 Aug 2019	AEM, OJ, CFC, ECC, MLB	192	55	34	69
Masisea island	23–24 Aug 2019	AEM, OJ, CFC, ECC, MLB	94	17	12	22
Headwater islands	6–7, 9–12 Oct 2015; 16 Aug 2018; 1, 26–27 Sep 2018	AEM, OJ, CFC, AEH, GFS	150	118	35	31
Middle Ucayali River	6, 11 Dec 2015; 5, 8–9, 19, 23–24 Aug 2019	AEM, OJ, CFC, ECC, MLB, MGH	72	0	0	10
Headwater rivers	6–7, 12 Oct 2015; 14, 16, 24–25 Aug 2018; 10, 24 Sep 2018; 1–2 Oct 2018	AEM, OJ, CFC, AEH, GFS	79	0	0	28
Preferida	31 Jul 2019; 1, 5–20 Aug 2019	AEM, OJ, CFC, ECC, MLB, MGH	298	310	108	127
Laguna Juanacha	18, 21–22 Aug 2019	AEM, OJ, CFC, ECC, MLB, MGH	197	25	11	76
Quebrada Caco	6–9 Dec 2015	MGH, EB	191	22	14	0
Breu	13–17 Dec 2015	MGH, EB	248	39	25	0
Atalaya area	6, 8 Oct 2015	AEM, OJ, GFS	102	0	0	10
Oventeni area	2–4 Jul 2015; 18–22 Sep 2015; 1–4 Oct 2015	AEM, OJ, GFS	221	101	62	26
Sira foothills	9, 13 Oct 2015	AEM, OJ, GFS	133	13	8	30
Cushireni	25 Sep–1 Oct 2018	AEM, CFC, AEH	77	48	22	7
Mayapo	27 Sep–1 Oct 2018	AEM, CFC, AEH	160	20	16	13
Serjali	13–14 Sep 2018	AEM, CFC, AEH	82	20	14	2
Sepa	11–13 Sep 2018	AEM, CFC, AEH	88	19	13	7
Nuevo San Martín	16–22 Sep 2018	AEM, CFC, AEH	171	38	18	11
~30 km NW Sepahua	2–9 Sep 2018	AEM, CFC, AEH	221	51	25	45
Sepahua	17–24, 29–31 Aug 2018	AEM, CFC, AEH	215	62	29	56
Camisea	25–29 Aug 2018	AEM, CFC, AEH	97	18	5	8

^a Audio recording and/or photograph in Macaulay Library.

the Ucayali near the city of Atalaya. We also sampled along the upper Juruá River at Breu and at elevations up to 1,100 m in the Cerros del Sira foothills and Oventeni. Here we describe each of these 22 sites, which are listed roughly from north to south, with a few adjustments to cluster sites of the same habitat (rivers and riverine islands) or same riverbank/interfluvium.

Yarinacocha—Although it is located just north of Pucallpa, we present our surveys from Yarinacocha (8.344°S, 74.571°W; ~140 m elevation) as a distinct locality due to its historical importance as a research locality (e.g., O'Neill and Pearson 1974). We used boats to survey the eastern half of the lake, and we also traversed the eastern edge of the lake and adjacent forest by foot.

Pucallpa area—This locality includes surveys from within the city itself, such as around the Hotel Ecológico Manish, FAP Captain David Abensur Rengifo International Airport, and the principal Plaza de Armas, but also sites well outside the city. These more distant sites included a dirt track ~17 km SW Pucallpa (8.504°S, 74.638°W; ~160 m elevation) and sightings within the Ucayali Region along the principal roads between Pucallpa and Campoverde and between Campoverde and Tournavista.

Tournavista area—Survey effort for this locality was concentrated along a 2 km stretch of dirt road 13 km west of Tournavista (8.94°S, 74.83°W; ~240 m elevation), a town on the west bank of the Pachitea River, which served as our base during




Figure 1. Map of study area in the central Amazon of Peru. The dotted arc passing by the localities of Masisea, Preferida, and Laguna Juanacha denotes the approximate route of the Ucayali River prior to the Ucayali avulsion (see Pärssinen et al. 1996).

fieldwork. Although this stretch of road provided access to some of the most extensive patches of *terra firme* forest in the area, the habitat was highly fragmented and interspersed with open agricultural areas. Forest clearing and road building were actively taking place in the immediate surroundings. This locality also includes observations from within the Huánuco Region along the road between Tourmavista and Campoverde. The habitat along this road was open agriculture with scattered trees and occasional small roadside marshes, several of which we stopped to survey.

Masisea island—This locality refers to a 215 ha island in the Ucayali River ~10 km north of the town of Masisea. The island was near the east bank, from which it was mostly separated by a narrow channel (satellite imagery revealed that the southern end of the island had recently connected to the mainland shore). Our surveys covered the grassy perimeter and small patches of *Cecropia* (Urticaceae) on the eastern side of the island (8.514°S, 74.309°W). The thick vegetation (mostly grasses) and lack of trails made for difficult work conditions and prevented access to some of the larger patches of *Cecropia*.

Headwater islands—For our surveys of islands in the Tambo, Urubamba, and upper Ucayali Rivers, we targeted those not visited by Harvey et al. (2014b). We visited 2 islands downriver of Atalaya on the Ucayali River: Isla Cetical (10.642°S, 73.841°W) and Isla del Gallo (10.675°S, 73.797°W). On the Urubamba River we visited 3 islands to the east of Atalaya (Isla Escondida, 10.737°S, 73.606°W; Isla Pukani, 10.692°S, 73.543°W; and an unnamed island, 10.721°S, 73.566°W) and an island farther upriver near Sepahua (Isla Pajuya, 11.066°S, 73.105°W). We also surveyed the gravel beaches of a complex set of small islands in the Tambo River (11.164°S, 73.710°W) but did not have a chance to survey the island interiors. Three of the islands visited (Cetical, del Gallo, and Escondida) were moderate in size (150–400 ha) and contained typical riverine island scrub habitat: a mix of mature *Cecropia* forest at the island center, with scrub of decreasing stature toward the periphery dominated by *Gynerium* (Poaceae) cane and *Tessaria* (Asteraceae) shrubs. Isla Pajuya was smaller in size (35 ha) but with habitat similar to that of the islands described above. Isla Pukani was larger in size (290 ha), and

its forest was taller, taking on a more *várzea*-like stature. Finally, the unnamed island in the Urubamba River was small (25 ha) and little more than a gravel bar with scattered forbs and grasses. Both Isla Pukani and Isla del Gallo were close to the riverbank and appeared to be merging with the shoreline. Every surveyed island larger than 50 ha contained some amount of habitat clearing for agricultural use.

Middle Ucayali River—To access other field sites we regularly traveled along the middle Ucayali River in boats, and we took advantage of these trips to document birds along the riverbanks and gravel bars. All surveys for this locality were along the stretch of river between Pucallpa and Masisea, except for one between the mouth of the Pachitea River and Quebrada Caco (see locality below).

Headwater rivers—Our extensive boat travel along the upper Ucayali River around Atalaya, Urubamba River (from Atalaya to Camisea), and Tambo River (between Atalaya and Puerto Ocopa) provided opportunities to survey birds along the riverbanks and gravel bars. In this locality, we also have included sightings from the Atalaya waterfront.

Preferida—This locality includes surveys at an established camp on land of the Shipibo community of Preferida de Charasmaná (which locals referred to simply as “Preferida”), 10.8 km southeast of Masisea (8.685°S, 74.250°W; 150 m elevation). It also includes a few surveys in the town of Masisea and along the roads in the immediate area. This site is in the eastern portion of the ~2,000 km² land area passively transported from the west to the east bank of the Ucayali River by the Ucayali avulsion. The primary habitat was tall, relatively intact *terra firme*-like forest, but included some forest in swampy terrain, agricultural fields, and edge habitats.

Laguna Juanacha—We launched a boat from the lakeside town (*caserío*) of Charasmaná (8.693°S, 74.206°W; not to be confused with the community of Preferida de Charasmaná detailed above) to survey the northeastern end of Laguna Juanacha and surrounding forest (8.694°S, 74.201°W; 155 m elevation). Surveys for this locality also involved a short trip along the Tamaya River to Cocha Larga (8.679°S, 74.191°W), an oxbow lake 2 km northeast of Charasmaná. In contrast to Cocha Larga, Laguna Juanacha is a blocked valley lake

formed during the Ucayali avulsion (Neller et al. 1992). The habitat in the area surrounding Laguna Juanacha and nearby Cocha Larga was largely *igapó* forest. We visited during the dry season, but high-water markings suggest that much of this forest is flooded to a depth of ~1.5–2 m for part of the year. Near Laguna Juanacha the understory was particularly open, but near Cocha Larga the understory was dense and viny. Cocha Larga was marsh-like with little open water, whereas Laguna Juanacha held extensive open water. Large mats of aquatic vegetation along parts of Laguna Juanacha were dominated by *Polygonum* sp. (Polygonaceae) or a close relative, and the forested margins of the lake were of particularly short stature (~5–7 m).

Quebrada Caco—Our surveys at this locality (9.403°S, 74.163°W; 160–170 m elevation) in 2015 covered the lower stretch of this blackwater creek immediately upstream from the town of Pueblo Nuevo del Caco. Habitats surveyed included a large area of seasonally flooded scrubby swamp and grasslands and nearby seasonally flooded forest (*igapó*).

Breu—This locality lies near the Brazil border and is the only Peruvian town in the watershed of the Juruá River (9.534°S, 72.762°W; 240 m elevation). We conducted surveys near town on both sides of the Juruá River, covering habitats including floodplain forest (*várzea*) with some extensive swamps and *terra firme* forest on hilly terrain containing many clearings and viny tangles.

Atalaya area—Our surveys for this locality were limited to a marsh 10 km south of Atalaya (10.821°S, 73.760°W) and nearby low foothill forest at ~500 m elevation (10.820°S, 73.771°W) on 8 October 2015. We also include incidental records within the city of Atalaya.

Oventeni—We surveyed a marsh, *pajonales* (grasslands), and forest patches below 1,100 m elevation around Oventeni (10.759°S, 74.221°W; 1,000 m elevation), the principal town in the Gran Pajonal, and the Ashéninka community of Kirahuanero (10.699°S, 74.218°W; 6.5 km to the north). For most of our time in the Oventeni area we were based at a camp on the eastern edge of Kirahuanero.

Sira foothills—This locality encompasses numerous sites (700–1,100 m elevation) along the primary dirt road between Puerto Ocopa and Atalaya. We stopped regularly along this road to conduct short surveys and to collect specimens but

did not establish any camps. Stops included a logging track in humid forest ~20 km southwest of Atalaya (10.915°S, 73.905°W; 880 m elevation), scrubby dry forest 22.3 km southeast of Oventeni (10.904°S, 74.082°W; 800 m elevation), a *pajonal* 19.4 km southeast of Oventeni (10.894°S, 74.104°W; 1,060 m elevation), and humid forest 15.4 km south of Oventeni (10.899°S, 74.195°W; 780 m elevation).

Cushireni—We based our work along the Tambo River from our camp (11.153°S, 73.724°W) on the west bank ~5 km downriver from the Ashéninka community of Cushireni (11.197°S, 73.708°W). We boated most days to the community itself to access their well-maintained trails. The primary habitat was tall foothill forest, which we surveyed up to 450 m elevation. We also surveyed edge habitats around agricultural clearings.

Mayapo—We conducted surveys in a variety of habitats around the Ashéninka community of Mayapo (11.171°S, 73.694°W; 275 m elevation), including transitional *várzea/terra firme* forest, young second-growth and edge habitats around clearings, and small tracts of *Guadua* bamboo. To survey mature *terra firme* forest (400 m elevation), AEM also dedicated 2 d along the well-maintained trail connecting the communities of Mayapo and Nazareth and spent a single night on the trail ~9 km east of Mayapo (11.154°S, 73.614°W).

Serjali—We established a camp near this small town on the west (left) bank of the Urubamba River ~25 km east of Atalaya (10.715°S, 73.525°W). We focused our surveys primarily on the transitional *várzea/terra firme* forest ~2 km inland from the river (280 m elevation).

Sepa—We established a camp on the west bank of the Sepa River about 2 km upstream from the mouth (10.835°S, 73.296°W). The river was shallow in many places (~0.5 m) during our mid-September visit, and it was only by unloading and frequently pushing our boat that we reached our field site. We surveyed *várzea* and hilly *terra firme* forests as well as clearing edges (230–365 m elevation).

Nuevo San Martín—Our camp along the Inuya River was based on the south bank in the Amahuaca community of Nuevo San Martín (10.587°S, 73.173°W; 210 m elevation). We surveyed pristine *terra firme* habitats on both the south and north banks of the Inuya River, with the

north bank being notably hilly (up to 250 m elevation). We also surveyed several small tracts of bamboo on both banks, edge habitats surrounding the community, and the river margins.

~30 km NW Sepahua—Our camp at this locality (10.911°S, 73.175°W; 250 m elevation) was located on the east bank of the Urubamba River on the edge of extensive *várzea* forest. We established a trail northeast to a small oxbow lake (10.903°S, 73.168°W) and continued until reaching logging trails that allowed for easier hiking. As we hiked into slightly higher elevations, we encountered vast tracts of *Guadua* bamboo. We established a satellite camp (10.889°S, 73.161°W), where we stayed for 6 nights, in a large bamboo tract ~3 km northeast of our principal camp. From the satellite camp we also accessed hilly *terra firme* forest farther north (10.884°S, 73.163°W; 300 m elevation).

Sepahua—We established 3 different camps near the town of Sepahua (11.144°S, 73.044°W) in 2018: first at ~9 km southeast of Sepahua on the east bank of the Urubamba River (11.193°S, 72.979°W; 17–20 August; 270 m elevation); second at 10 km east-southeast of Sepahua on the south bank of the Sepahua River (11.165°S, 72.957°W; 20–24 August; 290 m elevation); and third at 10.5 km east-southeast of Sepahua on the north bank of the Sepahua River (11.165°S, 72.952°W; 29–31 August; 290 m elevation). The terrain at the first camp rose steeply from the river and allowed us to quickly reach hilly *terra firme* forest up to just over 400 m elevation. Treefall gaps were common, perhaps related to the steepness of the terrain, which fostered thick undergrowth and viny tangles. There was also an overgrown *chacra* with similarly dense habitat. The second camp was situated in relatively intact habitat (some evidence of selective logging) with easy access to rolling *terra firme* forest. The nearby third site was at the edge of an overgrown *chacra*, but upon hiking north from camp this quickly transitioned to hilly *terra firme* forest, which we surveyed up to 330 m elevation.

Camisea—Our Camisea camp (11.726°S, 72.921°W) was located on the north (right) bank of the Camisea River ~4 km southeast of the community of Camisea. From this base, we focused our surveys in hilly *terra firme* forest on both banks of the river (up to 400 m elevation on

the north bank) but also surveyed birds in the river-edge forest around camp.

Fieldwork

Daily fieldwork consisted of a combination of mist-netting throughout the day and audiovisual surveys concentrated in the mornings and evenings. To collect specimens, we deployed mist nets and used shotguns. We also obtained sound recordings and photos of as many species as possible. We prepared study skins for most specimens and prepared a smaller number as skeletons. For all specimens, we flash-froze pectoral muscle, heart, and liver tissues in liquid nitrogen, with additional muscle and stomach content samples preserved in ethanol. For at least 1 individual per species, we also preserved the lower intestine in liquid nitrogen for gut microbiota studies. We submitted complete lists of species observed, often several per day, to the Avian Knowledge Network through the eBird portal (Sullivan et al. 2009), which is freely accessible online. We uploaded our bird photographs and audio recordings to the Macaulay Library (<https://www.macaulaylibrary.org>) through the eBird portal. ML numbers in the below text refer to supporting media archived online at the Macaulay Library and XC numbers refer to supporting audio recordings archived online at <https://www.xeno-canto.org>. LSUMZ numbers refer to catalogued specimens in the bird collection at the LSUMNS (formerly the Louisiana State University Museum of Zoology). We follow Remsen et al. (2021) for English and scientific names of birds.

Results

We detected 657 species and collected 980 specimens of 294 species (Table 1; see Supplemental Table S1 for complete species list). We evenly divided specimen collections between CORBIDI and LSUMNS, and both institutions received tissue samples for all individuals. Below we describe avian communities associated with major habitat types and summarize novel biogeographic patterns. More detailed accounts for notable species are included in the Supplemental Appendix.

Avifauna by habitat types

Birds of upland/terra firme forest—Our most intensive surveys in *terra firme* forest occurred around the Preferida camp within the area of land transported by the Ucayali avulsion. The forest here was swampier and had a thicker understory than is typical of *terra firme*. We detected species typical of *terra firme* forest such as White-throated Tinamou (*Tinamus guttatus*), Black-bellied Cuckoo (*Piaya melanogaster*), Blue-cheeked Jacamar (*Galbula cyanicollis*), Paradise Jacamar (*Galbula dea*), Spot-winged Antbird (*Myrmelastes leucostigma*), Blue-crowned Manakin (*Lepidothrix coronata*), White-crested Spadebill (*Platyrinchus platyrhynchos*), and Brownish Twistwing (*Cnipodectes subbrunneus*) but also some species typical of *várzea* or transitional forests such as Cinereous Tinamou (*Crypturellus cinereus*), Spotted Puffbird (*Bucco tamatia*), Rufous-necked Puffbird (*Malacoptila rufa*), White-shouldered Antbird (*Akletos melanoceps*), Band-tailed Manakin (*Pipra fasciicauda*), and Gray-crowned Flycatcher (*Tolmomyias poliocephalus*). Interestingly, this site also harbored some local species typical of forest on nutrient-poor soil such as Brown-banded Puffbird (*Notharchus ordii*) and Citron-bellied Attila (*Attila citriniventris*).

Other *terra firme* sites sampled on both banks of the Ucayali, Tambo, and Urubamba rivers overlapped with Preferida in avian species composition, but generally had more open understory and hilly terrain that reached higher elevations relative to the floodplain. This was likely responsible for the detection of additional *terra firme* bird species (Supplemental Table S1). In the foothills of the Sira these included notable species such as Foothill Elaenia (*Myiopagis olallai*) and Blackish Pewee (*Contopus nigrescens*), both of which are poorly known in central Peru (see Supplemental Appendix). In contrast to most *terra firme* sites, forest at our principal Tournavista site west of the Ucayali River was severely fragmented. Bird diversity was still high, and we encountered most of the common species detected in *terra firme* sites elsewhere. We failed, however, to detect some expected taxa including anthrushes (*Formicarius* spp.), Long-tailed Woodcreeper (*Deconychura longicauda*), White-chinned Woodcreeper (*Dendrocincla merula*), Wedge-billed Woodcreeper (*Glyphorynchus spirurus*), foliage-gleaners (*Phil-*

ydor and *Automolus* spp.), and Pipridae spp. (except for one male Blue-crowned Manakin). Further work is needed to identify whether these absences from the avian community are associated with fragmentation, which can lead to reduced diversity and even extirpation of mixed-species flocks (Rutt et al. 2020), or are due to limited sampling.

Birds of floodplain/várzea forest—Our surveys of *várzea* forest were restricted mostly to the banks of the Urubamba River. We surveyed this habitat most thoroughly near our camp ~30 km northwest of Sepahua, but our mist-netting and general survey efforts were limited in this habitat compared to those in *terra firme* forest at this and other sites. Still, we detected many species typical of *várzea* forest (Supplemental Table S1). We also detected in this habitat several species typical of *terra firme* forest, notably 2 Fulvous-chinned Nunlets (*Nonnula sclateri*) captured in mist nets a few meters up-bank from the river at our camp ~30 km northwest of Sepahua.

Birds of blackwater flooded forests/igapó—Our surveys at Quebrada Caco, Laguna Juanacha, and Cocha Larga are the most extensive conducted in blackwater habitats in the central Amazon of Peru. The previous lack of attention paid to this habitat is highlighted by our detection of 5 new species for the Ucayali Region in just 3 d of fieldwork: Band-tailed Nighthawk (*Nyctiprogne leucopyga*), Green-tailed Goldenthrout (*Polytmus theresiae*), Yellow-chinned Spinetail (*Certhiaxis cinnamomeus*), Amazonian Black-Tyrant (*Knipolegus poecilocercus*), and Gray-chested Greenlet (*Hylophilus semicinerus*). The avifauna at these sites resembles that noted in blackwater habitats at the Pacaya-Samiria National Reserve in the Loreto Region (Begazo and Valqui 1998). Characteristic forest species included Scarlet-crowned Barbet (*Capito aurovirens*), Plain-breasted Piculet (*Picumnus castelnaui*), Black-crested Antshrike (*Sakesphorus canadensis*), Amazonian Antshrike (*Thamnophilus amazonicus*), Stripe-chested Antwren (*Myrmotherula longicauda*), Black-tailed Antbird (*Myrmoborus melanurus*; at Cocha Larga), Yellow-crowned Elaenia (*Myiopagis flavivertex*), and Cinnamon Attila (*Attila cinnamomeus*). Green-tailed Golden-throat and Gray-chested Greenlet were common components of the avifauna in scrubby lakeside forest at Laguna Juanacha, and Band-tailed Nighthawk was abundant in the hour before dawn,

singing from lakeside vegetation and foraging over the lake. Additional species in edge and scrub habitats included Cinereous Becard (*Pachyrampus rufus*) and Dot-backed Antbird (*Hylophylax punctulatus*) at Quebrada Caco and Amazonian Black-Tyrant at Laguna Juanacha. Dense floating stands of aquatic *Polygonum* vegetation seemed to be the preferred habitat for Yellow-chinned Spine-tail, and Pied Water-Tyrant (*Fluvicola pica*) was also present at lake margins.

Birds of Guadua bamboo—During fieldwork in 2018 along the headwaters of the Ucayali, we detected most of Kratter's (1997) avian specialists of *Guadua* bamboo habitats of the southwestern Amazon Basin. The composition of the bamboo bird communities surveyed varied greatly depending on the extent of bamboo present. The largest and most mature bamboo tract was ~30 km northwest of Sepahua (10.889°S, 73.161°W; 300 m elevation) and ~3 km northeast of our camp on the bank of the Urubamba River. This tract extended for ~1 km along a trail and was composed of dense, ~10 m tall bamboo, bordered by *terra firme* forest to the northeast and cleared pasture to the southwest. Obligate and near-obligate bamboo specialists (Kratter 1997, Lebbin 2013) detected at this site included Pavonine Cuckoo (*Dromococcyx pavoninus*), Yellow-billed Nunbird (*Monasa flavirostris*), Rufous-headed Woodpecker (*Celeus spectabilis*), Striated Antbird (*Drymophila devillei*), Manu Antbird (*Cercomacra manu*), White-lined Antbird (*Myrmoborus lophotes*), Goeldi's Antbird (*Akletos goeldii*), Dusky-cheeked Foliage-Gleaner (*Anabazenops dorsalis*), Peruvian Recurvebill (*Syndactyla ucayalae*), Brown-rumped Foliage-Gleaner (*Automolus melanopezus*), Flammulated Pygmy-Tyrant (*Hemitriccus flammulatus*), White-cheeked Tody-Flycatcher (*Poecilatriccus albifacies*), and Large-headed (*Ramphotrigon megacephalum*) and Dusky-tailed (*R. fuscicauda*) flatbills. Several facultative bamboo-associated species (Kratter 1997, Lebbin 2013) were also present in the area, including Chestnut-capped Puffbird (*Bucco macrodactylus*), Rufous-breasted Piculet (*Picumnus rufiventris*), Red-billed Scythebill (*Campylorhamphus trochilirostris*), and Long-crested Pygmy-Tyrant (*Lophotriccus eulophotes*). Notably, this was the only site where we detected Pavonine Cuckoo, Yellow-billed Nunbird, and Peruvian Recurvebill.

The tracts of bamboo we surveyed elsewhere were smaller, but each generally contained at least

a few bamboo specialists. Around Nuevo San Martín, the bamboo occurred on both sides of the Inuya River (10.599°S, 73.171°W and 10.573°S, 73.190°W), and several specialists recorded here represent some of the northernmost records for their species in Peru (e.g., Manu Antbird and White-cheeked Tody-Flycatcher). The bamboo at Mayapo was part of a matrix of habitat that included *terra firme*, cleared cultivated land, and viny tangles near water. We found many bamboo specialists here (see Supplemental Table S1), including 2 species not detected elsewhere: Ornate Antwren (*Epinecrophylla ornata*) and the recently described Rufous Twistwing (*Cnipodectes super-rufus*; Lane et al. 2007). At our first Sepahua locality, we collected a pair of the typically bamboo-associated Bamboo Antshrike (*Cymbilaimus sanctaemariae*) in a matrix of *terra firme* and viny tangles but failed to detect this species in bamboo at any of our study sites.

Birds of riverine islands—The bird communities of the 5 typical riverine islands (Masisea, del Gallo, Cetical, Escondida, and Pajuya) were similar between islands and were dominated by a combination of obligate riverine island species—of which we detected 14 of the 18 species found in western Amazonia (Rosenberg 1990)—and species typical of open country and disturbed Amazonian habitats (Supplemental Table S1). We detected all of the island specialists found by Harvey et al. (2014b) and added Brownish Elaenia (*Elaenia pelzelni*) and Bicolored Conebill (*Conirostrum bicolor*), both of which were seen only on the Masisea island. Riverine island specialists detected at Isla Pajuya (near Sepahua) were Olive-spotted Hummingbird (*Leucippus chlorocercus*), Parker's Spinetail (*Cranioleuca vulpecula*), Castelnau's Antshrike (*Thamnophilus cryptoleucus*), and Black-and-white Antbird (*Myrmochanes hemileucus*), the latter 2 species representing the southernmost records in Peru. The avifauna of Isla Pukani had a distinctly *várzea*-like component, with species such as Silvered Antbird (*Sclateria naevia*) and Amazonian Antpitta (*Myrmothera berlepschi*). Little more than an exposed gravel bar, the small unnamed island in the Urubamba River was depauperate, but the riverine island specialists Lesser Hornero (*Furnarius minor*) and River Tyrannulet (*Serpophaga hypoleuca*) were both present despite the extremely young age of the island. In addition to the riverine

island specialists, we detected a variety of boreal migrant shorebird species, both austral and boreal migrant passerine species, and 3 intra-tropical migrant *Sporophila* species on riverine islands.

Birds of open country and wetlands—We conducted occasional avifaunal surveys of open agricultural habitats, which are rapidly increasing in extent in the central Peruvian Amazon, particularly in the greater Pucallpa area (Bax et al. 2016, Vijay et al. 2018). In addition to a variety of expected open country species (Supplemental Table S1), we observed Crested Caracara (*Cara-cara plancus*), a recently arrived species in the area (Piana et al. 2012; see Supplemental Appendix), on several occasions. Marshes within agricultural areas along the dirt road between Tournavista and Campoverde held a variety of waterbirds including less expected species such as Snail Kite (*Rostrhamus sociabilis*) and Pied Water-Tyrant (*Fluvicola pica*). Naturally occurring open habitats and wetlands around Lago Yarinacocha provide an important foraging area for birds near the city of Pucallpa (Supplemental Table S1). Boreal-breeding migratory shorebirds are poorly surveyed in the central Peruvian Amazon, but our observations suggest Lago Yarinacocha may be an important stopover site. On 20 August 2019 we encountered ~700 Pectoral Sandpipers (*Calidris melanotos*) and ~450 Lesser Yellowlegs (*Tringa flavipes*) at Yarinacocha, and on 31 August 2019 we encountered ~250 Stilt Sandpipers (*Calidris himantopus*) and ~800 Lesser Yellowlegs. At Yarinacocha we also documented American Golden-Plover (*Pluvialis dominica*), Buff-breasted Sandpiper (*Calidris subruficollis*), and Wilson's Phalarope (*Phalaropus tricolor*), and along the Ucayali River we noted Black-bellied Whistling-Duck (*Dendrocygna autumnalis*), Brazilian Teal (*Amazonetta brasiliensis*), Franklin's Gull (*Leucophaeus pipixcan*), and Puna Ibis (*Plegadis ridgwayi*), all species poorly known in the Peruvian Amazon (see Supplemental Appendix).

We conducted additional surveys of open country in the *pajonales* around Oventeni and on the Puerto Ocopa–Atalaya Road (~1,000–1,100 m), areas previously surveyed by Harvey et al. (2011). We found many species typical of open country in the lowlands, but we also detected some species only in this area including Spot-tailed Nightjar (*Hydropsalis maculicaudus*), Ocellated Crane (*Micropygia schomburgkii*), Russet-

crowned Crane (*Anurolimnas viridis*), Rusty-backed Antwren (*Formicivora rufa*), Cinereous-breasted Spinetail (*Synallaxis hypospodia*), Yellow-bellied Elaenia (*Elaenia flavogaster*), Lesser Elaenia (*Elaenia chiriquensis*), Yellow-browed Tyrant (*Satrapa icterophrys*; austral migrant), Black-faced Tanager (*Schistochlamys melanopsis*), Wedge-tailed Grass-Finch (*Emberizoides herbicola*), and Plumbeous Seedeater (*Sporophila plumbea*).

Species turnover across rivers

We found that species distributions nearly all conformed to previous findings for river-limited taxa in the central Peruvian Amazon (Harvey et al. 2014b). Harvey et al. (2014b) noted 9 species pairs with distributions limited by the middle Ucayali, and our surveys largely support those findings. Even at Preferida, which was on the west bank of the Ucayali River until ~230–250 years ago (Pärssinen et al. 1996), we found almost entirely east-bank taxa, including Blue-cheeked Jacamar (*Galbula cyanicollis*), Fulvous-chinned Nunlet (*Nonnula sclateri*), Bar-breasted Piculet (*Picumnus aurifrons*), White-bellied Parrot (*Pionites leucogaster*), White-throated Antbird (*Oneillornis salvini*), White-bellied Tody-Tyrant (*Hemitriccus griseipectus*), and Moustached Wren (*Phlegopedius genibarbis*). However, we did find 2 apparent instances of west-bank taxa at Preferida: a specimen of Rufous-rumped Foliage-Gleaner (*Philydor erythrocerum*; LSUMZ 228870) that closely resembles the darker brown *P. e. subfulvum* from west of the Ucayali River rather than the expected *P. e. lyra* and a specimen of Double-banded Pygmy-Tyrant (*Lophotriccus vitosus*; LSUMZ 228926) that shows gray crest margins similar to *L. v. vitosus* from west of the river rather than the buffy or olive crest margins of *L. v. congener*, the expected taxon east of the river. In addition, we noted 2 striking cases of apparent hybridization in the area of the avulsion.

First, we found evidence of introgressive hybridization between White-bellied Parrot and Black-headed Parrot (*Pionites melanocephalus*) along the middle Ucayali River (Fig. 2a). Haffer (1977) had previously noted several specimens with characteristics intermediate between these species in eastern Peru: White-bellied Parrots with different combinations of scattered black crown



Figure 2. Specimens of 2 species pairs bounded by the Ucayali River show evidence of introgression across the river. In each panel, a blue line indicates the divide between west/left bank birds from Tournavista and east/right bank birds from Preferida, which is located within the patch of *terra firme* forest that was transported from the west bank to the east bank during the Ucayali avulsion. (a) *Pionites* parrots: Black-headed Parrot (*P. melanocephalus*) is expected on the west bank and White-bellied Parrot (*P. leucogaster*) on the east bank. The 3 west bank individuals (left to right: LSUMZ 228716, 228718, 228713) show largely *P. melanocephalus* phenotypes but with varying degrees of *leucogaster*-like traits: LSUMZ 228716 appears close to the parental type, but the other 2 show orange feathering in the forehead and crown and the amount of green in the lores is reduced, especially in LSUMZ 228713. Six individuals from the east bank at Preferida are shown (left to right: LSUMZ 228725, 228726, 228720, 228727, 228729, 228730). One east bank individual, LSUMZ 228725 (fourth from left overall), shows a *melanocephalus*-like phenotype, despite being on the “wrong” side of the Ucayali River. Like *P. melanocephalus* it has a fully black crown, bill, and orbital skin, but unlike *P. melanocephalus* it has very little green in the lores. Among the other more *leucogaster*-like individuals, there is substantial variation in the presence of *melanocephalus*-like traits, such as black feathers in the crown, dark coloring of the bill and orbital skin, and green feathers in the lores. (b) *Pheugopedius* wrens: Coraya Wren (*P. coraya*) is expected on the west bank and Moustached Wren (*P. genibarbis*) on the east bank. The individual from the west bank (LSUMZ 228997) is typical for *P. coraya*, with dark throughout the auriculars, malar, and lateral throat stripe and rich chestnut coloration on the flanks and belly. The 4 individuals from Preferida on the east bank (left to right: LSUMZ 228994, 228992, 228996, 228988) show variation in their facial patterns, including in the degree of white streaking in the auriculars and the extent to which a white malar divides the auriculars from the dark lateral throat stripe. The color of the flanks also varies, but interestingly does not seem to correspond with the face pattern: the individual with the most *coraya*-like face pattern (LSUMZ 228994) has among the lightest and least extensive chestnut coloration on the underparts.

feathers, dark orbital skin, and dusky bill east of the Ucayali River and a Black-headed Parrot (AMNH 237774) with rusty orange on the forecrown whose date of collection (5 August

1927) suggests it was east of the Ucayali River (see Wiley 2010). Some of the specimens we collected showed similar morphological intermed-
diacy. In our series of 10 adult White-bellied

Parrot-like individuals from Preferida, we found that 6 had variable but substantial amounts of black feathering on the hind-crown and 4 had essentially pure, rusty orange crowns. Three individuals from this series displayed particularly dark orbital skin and 6 had noticeably dusky coloration on their bills, but these traits were quite variable. Some individuals (e.g., LSUMZ 228727 and 228729) also had a few scattered green feathers in the lores. Also at our Preferida camp, we collected an 11th *Pionites* that appeared more like an adult Black-headed Parrot (LSUMZ 228725) with a fully black crown, bill, and orbital skin, yet with very limited greenish in the lores. In our series of 7 adult Black-headed Parrot-like individuals from Tournavista, we also observed variable amounts of black feathering on the crown, with several individuals having a fully black crown and several showing extensive rusty orange on the forecrown. This suggests introgression with White-bellied Parrot, a species that remains undocumented west of the middle Ucayali River. A recent photograph in an eBird report indicates presence of some White-bellied Parrot-like individuals west of the lower Ucayali (Rowland 2018; ML216708631), and Harvey et al. (2014b) documented White-bellied Parrot west of the upper Ucayali. Although we currently lack confirmatory genetic data, we suggest that the phenotypes of birds sampled here are indicative of substantial introgression not just around the Ucayali avulsion, but more widely in the middle Ucayali Valley. The observation that adult White-bellied Parrots can have scattered black feathers on the crown up to nearly 500 km southeast of the Amazon River in northeastern Pará, Brazil (Novaes 1981), may suggest either that introgression occurs far from zones of potential contact, or that phenotypic variation within White-bellied Parrot is complex. Further collections and high-quality photographs of *Pionites* are greatly needed to assess plumage variation in and away from contact zones.

Second, we found that Coraya Wren (*Pheugopedius coraya*) and Moustached Wren appear to hybridize across the middle Ucayali River (Fig. 2b). Previous studies suggested this might be the case: O'Neill and Pearson (1974) noted both species at Yarinacocha and Schulenberg et al. (2010) noted intergradation in plumage features of birds on the west bank of the Ucayali. At our Preferida site, on the east bank, we collected a

series of 9 *Pheugopedius* wrens that exhibit substantial variation in the extent of white streaking on the auriculars (mostly like Moustached Wren but in some individuals approaching the mostly black auriculars of the Coraya Wren phenotype). In this series, variation was also notable in the extent of the white malar stripe directly above (and which highlights) the black lateral throat stripe characteristic of Moustached Wren. In some cases, the white malar stripe was essentially absent or faint, which resulted in nearly continuous black feathering from the lateral throat stripe to the auriculars in several individuals. These intermediate plumage features in the Preferida series suggest hybridization. The single individual of Coraya Wren that we collected at Tournavista (LSUMZ 228997) shows the expected limited white streaking in the auriculars and lacks the white malar stripe. Vocal evidence also suggests hybridization between Coraya and Moustached wrens on the left bank of the Ucayali River. The populations of Coraya Wren in lowland Amazonia (*P. c. amazonicus*) and foothills of San Martín (*P. c. albiventris*) west of the Ucayali River and south of the Marañón River give the *djeer djeer* call (e.g., ML190519 and P. Boesman recording XC230995) that is typical of Moustached Wren and is lacking in Coraya Wren populations (e.g., *P. c. griseiventris* and *P. c. coraya*) north of the Marañón and Amazon Rivers and farther east (DFL, pers. obs.). This shared call between taxa across the Ucayali River seems most plausibly explained by a scenario of extensive gene flow, although, because call learning has been documented in oscine passerines (Löhr 1963, Bertram 1970, Munding 1979, Zann 1985), it is conceivable that this pattern is due to a shared and learned dialect.

Notable distributional records

We documented first regional records of 28 bird species across the Ucayali (18), Huánuco (3), and Junín (7) regions (Table 2) and numerous other geographic range extensions with implications for patterns of avian diversity and taxonomy (see Supplemental Appendix). Many of these notable records derive from our sampling of poorly known habitats in the area, particularly blackwater wetland systems, *Guadua* bamboo tracts, and riverine islands. Both blackwater systems and

Table 2. Summary of first regional records in the central Amazon of Peru in 2015, 2018, and 2019. Abbreviations: S = specimen, M = media (audio recording and/or photograph in Macaulay Library), X = only detected visually and/or aurally.

Common name	Scientific name	Region	Habitat type	Documentation
Band-tailed Nighthawk	<i>Nyctiprogne leucopyga</i>	Ucayali	Blackwater wetland	S, M
White-chinned Swift	<i>Cypseloides cryptus</i>	Ucayali, Huánuco	Aerial	M
Green-tailed Goldenthrout	<i>Polytmus theresiae</i>	Ucayali	Blackwater wetland	S, M
Wire-crested Thorntail	<i>Discosura popelairii</i>	Ucayali	Foothill forest	X
Semipalmated Plover	<i>Charadrius semipalmatus</i>	Ucayali	Riverine marsh/beach	X
Killdeer	<i>Charadrius vociferus</i>	Ucayali	Riverine marsh/beach	X
Wilson's Phalarope	<i>Phalaropus tricolor</i>	Ucayali	Riverine marsh/beach	S, M
Franklin's Gull	<i>Leucophaeus pipixcan</i>	Ucayali	Ucayali River	X
Bamboo Antshrike	<i>Cymbilaemus sanctaemariae</i>	Ucayali	<i>Terra firme</i> forest	S, M
Yellow-breasted Antwren	<i>Herpsilochmus axillaris</i>	Ucayali	Foothill forest	S, M
Dot-backed Antbird	<i>Hylophylax punctulatus</i>	Ucayali	Blackwater wetland	S
Yellow-chinned Spinetail	<i>Certhiaxis cinnamomeus</i>	Ucayali	Blackwater wetland, river island	S, M
Brownish Elaenia	<i>Elaenia pelzelni</i>	Ucayali	River island	S, M
Torrent Tyrannulet	<i>Serpophaga cinerea</i>	Ucayali	Foothill stream	X
Blackish Pewee	<i>Contopus nigrescens</i>	Ucayali	Foothill forest	S, M
Amazonian Black-Tyrant	<i>Knipolegus poecilocercus</i>	Ucayali	Blackwater wetland	S
Gray-chested Greenlet	<i>Hylophilus semicinereus</i>	Ucayali	Blackwater wetland	S, M
Southern Martin	<i>Progne elegans</i>	Ucayali	Aerial	M
Needle-billed Hermit	<i>Phaethornis philippii</i>	Huánuco	<i>Terra firme</i> forest	X
Variiegated Flycatcher	<i>Empidonomus varius</i>	Huánuco	<i>Terra firme</i> forest edge	M
Black-and-white Hawk-Eagle	<i>Spizaetus melanoleucus</i>	Junín	Foothill forest	M
Undulated Antshrike	<i>Frederickena unduliger</i>	Junín	<i>Terra firme</i> forest	M
Manu Antbird	<i>Cercomacra manu</i>	Junín	<i>Guadua</i> bamboo	S
Brown-rumped Foliage-gleaner	<i>Automolus melanopezus</i>	Junín	<i>Guadua</i> bamboo	S, M
Striolated Manakin	<i>Machaeropterus striolatus</i>	Junín	Foothill forest	S, M
Black-backed Tody-Flycatcher	<i>Poecilatriccus pulchellus</i>	Junín	Dense, second-growth forest	S
Shiny Cowbird	<i>Molothrus bonariensis</i>	Junín	Marsh	M

riverine islands have been extensively surveyed north of our study area along the lower Ucayali and upper Amazon rivers in the Loreto Region (Remsen and Parker 1983, Rosenberg 1990, Begazo and Valqui 1998, O'Shea et al. 2015, Socolar et al. 2018). Thus, the range extensions for birds in these habitats were all southward (~230–600 km). In contrast, our surveys in *Guadua* bamboo extended the known ranges of several specialists north and/or west from sites in the Cuzco and Madre de Dios regions (Kratter 1997, Parker et al. 1997, Lebbin 2007, Harvey et al. 2014a) and from Balta in far eastern Ucayali (O'Neill 1974, 2003).

Discussion

Our surveys in the central Amazon of Peru highlight a remarkably diverse avifauna and provide new information about how this diversity is partitioned across habitats and biogeographic

barriers. We found that, despite a long history of ornithological work in the area, visits to under-surveyed habitats revealed many species not previously detected. These results emphasize the importance of habitat specialization to Amazonian birds and the need for greater attention to the characteristics of surveyed habitats, including soil conditions, which have a key but underappreciated influence on avian distributions (Whitney and Alvarez 1998). Our results also refine the locations of range limits and contact zones with respect to rivers and the Cerros del Sira, pointing to additional geographic sampling priorities and revealing preliminary evidence of some biogeographic stability even in a dynamic landscape.

Our surveys were the most extensive to date in blackwater wetland systems, *Guadua* bamboo tracts, and riverine islands in the central Peruvian Amazon. However, these were still insufficient to develop a complete picture of the avian diversity of these habitats. Future surveys in blackwater

systems should target species such as Masked Duck (*Nomonyx dominicus*), Zigzag Heron (*Zenaidura macroura*), Rufous Potoo (*Nyctibius bracteatus*), Three-striped Flycatcher (*Conopias trivirgatus*), Sulphur-bellied Tyrant-Manakin (*Neopelma sulphureiventer*), and Orange-crowned Manakin (*Heterocercus aurantiivertex*) that are as yet unrecorded or are very poorly known in the area. The blackwater systems within the Área de Conservación Regional Imiría are even more extensive than the ones we visited and are a high priority for future surveys. The 4 obligate riverine island species still unrecorded in Peru's central Amazon are Ash-breasted Antbird (*Myrmoborus lugubris*), Leaden Antwren (*Myrmotherula assimilis*), Pale-billed Hornero (*Furnarius torridus*), and Pearly-breasted Conebill (*Conirostrum margaritae*), and they should be searched for especially on larger islands near Pucallpa. Zimmer's Woodcreeper (*Dendroplex kienerii*) is also found on riverine islands in northern Peru (Moncrieff et al. 2019) and should be looked for farther south. Similarly, *Guadua* bamboo tracts likely hold several species not yet documented or very poorly known in the area such as Ihering's Antwren (*Myrmotherula iheringi*), Slate-colored Seedeater (*Sporophila schistacea*), and the newly described Acre Tody-Tyrant (*Hemitriccus cohnhaffti*; Zimmer et al. 2013), a species only recently documented for Peru in the Madre de Dios Region (Harvey et al. 2014a). Stunted white-sand forests, a habitat type we did not visit during this fieldwork, remain a glaring gap for avifaunal surveys throughout most of the central Peruvian Amazon. The only surveys of this habitat in the area so far are along the Cohengua River (Harvey et al. 2014b), where records of Zimmer's Tody-Tyrant (*Hemitriccus minimus*), "Campina" Fuscous Flycatcher (*Cnemotriccus fuscatus duidae*), Yellow-throated Flycatcher (*Conopias parvus*), and Black Manakin (*Xenopipo atronitens*) remain over 200 km from the nearest records in Acre, Brazil (Whittaker and Oren 1999, Guilherme and Borges 2011).

In addition to its close association with habitat diversity, avian diversity in the study area is tied to landscape features, particularly the Ucayali River. We found that patterns of species replacement across the Ucayali were generally consistent with those observed previously (Harvey et al. 2014b), but the apparent presence of 2 subspecies on the "wrong" bank and hybridization in 2 species pairs

along the middle Ucayali suggests more complicated biogeographic histories in some species. Events such as the Ucayali avulsion of the 18th century have likely occurred regularly over the history of the river drainage (Pärssinen et al. 1996), passively transferring populations of birds to the "wrong" bank of the river. Despite the recent transfer of ~2,000 km² of land, including *terra firme* forest, to the east side of the river during the Ucayali avulsion, we found that most avian populations within this landmass morphologically matched the east bank species. This suggests that patterns of avian replacements around river barriers can recalibrate even after large-scale barrier adjustments. We suspect that the small populations of west bank birds that were passively transferred during the Ucayali avulsion quickly died out or were genetically swamped by the large populations of east bank birds. Future genetic work may provide additional clues about the fate of those populations and reveal insight into the impacts of landscape dynamics on populations.

Our knowledge of how habitat diversity and landscape features influence patterns of avian diversity helps to predict patterns in unsurveyed areas, but we emphasize that sometimes expected species simply do not occur at apparently suitable sites. Given our current limited data, it is often impossible to distinguish between true species absence and insufficient sampling. For example, our targeted surveys for Blue-crowned Manakins in *terra firme* at multiple localities in the Tambo-Urubamba interfluvium failed to detect any individuals, but we hesitate to suggest species absence from such a large area. A single specimen (AMNH 240618) collected by the Olalla brothers in 1927 from "Boca Rio Urubamba" appears to indicate presence of the species in the interfluvium, but modern verification is highly desirable. A second specimen from the same locality (AMNH 407206) appears to be from the problematic Harvey Bassler collection in which locality information is of dubious origin (see Wiley 2010). As we continue to conduct surveys, we ultimately hope to confidently identify distribution gaps and limits. In addition to species absence, species turnover away from rivers generally remains a mystery. For example, many pairs of related taxa including Red-billed/Rufous-vented Ground-Cuckoo (*Neomorphus pucheranii/N. geoffroyi*), Needle-billed/Straight-billed Hummingbird (*Phaethornis philip-*

pii/P. bourcierii), Rusty-breasted/Fulvous-chinned Nunlet (*Nonnula rubecula/N. sclateri*), Saturnine/Dusky-throated Antshrike (*Thamnomanes saturninus/T. ardesiacus*), White-eared/Purus Jacamar (*Galballyrhynchus leucotis/G. purusianus*), and Rufous-backed/Rio Madeira Stipplethroat (*Epinecrophylla haematonota/E. amazonica*) appear to turn over in continuous forest of central Peru east of the Ucayali River (Schulenberg et al. 2006, Del-Rio et al. 2021, Johnson et al. 2021), highlighting the need for finer resolution data to further investigate these biogeographic patterns.

Despite the high diversity and comparatively pristine condition of the central Peruvian Amazon, serious conservation issues plague the area. The most notable center of habitat loss is around the city of Pucallpa and along the Federico Basadre Highway (ruta nacional PE-18 C), a road established in the 1940s and a major determinant of regional deforestation patterns (Ichikawa et al. 2014, Bax et al. 2016). Perhaps foremost among the emerging threats to regional biodiversity is the rapid expansion of oil palm plantation monocultures in Peru (Vijay et al. 2018), which have been documented to hold depauperate bird communities around Pucallpa and farther abroad in Amazonia (Lees et al. 2015, Srinivas and Koh 2016). Although more focused in its effect, Pucallpa's thriving illegal animal trade also poses a conservation threat to numerous species of birds—primarily parrots (Gastañaga et al. 2011, Daut et al. 2015). Even now, the biggest asset for conservation of the central Peruvian Amazon is the very remoteness that has limited ornithological work. As this remoteness ebbs, efforts to avert catastrophic loss of avian diversity will benefit from a more complete understanding of regional avian diversity, habitat associations, and biogeography.

Acknowledgments

We thank RT Brumfield, JV Remsen, FH Sheldon, F Angulo, MA Plenge, TS Schulenberg, BM Whitney, reviewer JB Socolar, and one anonymous reviewer for discussion and comments that improved the manuscript. T Valqui, EP Derryberry, T Jackson, and C Quintori provided invaluable support with expedition logistics. M Rego and RH Wiley generously provided databases on relevant specimen collections. M Adams provided helpful photographs of specimens. We are also grateful to the Cornell Lab of Ornithology for loaning sound recording equipment. We would never have realized this fieldwork without the help of

local field assistants E Monsín, E Andrade, and J Hueroné and the trust and support of the Central Asháninka de Río Tambo (CART), Organización Distrital Indígena de Masisea (ORDIM), and the indigenous communities of Camisea, Cushireni, Kirahuanero, Mayapo, Nuevo San Martín, Preferida de Charasmaná, Sepahua, and Serjali.

Our fieldwork was funded by the Coypu Foundation, donations to the LSUMNS Tropical Bird Research Fund, and an American Ornithological Society grant to MGH. AEM, OJ, and AEH were supported during this work by National Science Foundation Graduate Research Fellowships. Any opinion, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation. AEH was also supported by a Louisiana Board of Regents Fellowship. No funding source had any influence on the content of the submitted or published manuscript, nor required approval of the final manuscript before publication. Specimen collection was authorized by the Servicio Nacional Forestal y de Fauna Silvestre (SERFOR) under permits 203-2015-SERFOR-DGGSPFFS and 222-2015-SERFOR-DGGSPFFS. Specimen collection was done under LSU's Institutional Animal Care and Use Committee (IACUC) protocol numbers 15-036 and 18-054.

Literature cited

- Álvarez A J, Metz MR, Fine PVA. 2013. Habitat specialization by birds in western Amazonian white-sand forests. *Biotropica*. 45:365–372.
- Angulo F, Novoa J, Lazo I. 2016. Aves del río La Novia en la cuenca del Purús, Ucayali, Perú [Birds of the La Novia River in the Purus drainage, Ucayali, Peru]. In: Luis M J, Germaná C, editors. *Diversidad biológica del sudeste de la Amazonía Peruana: avances en la investigación*. Lima (Peru): Ediciones Nova Print S.A.C. p. 122–128. Spanish.
- Balta AK. 2007. La avifauna del Río Yuruá [The avifauna of the Jurúa River]. *Boletín de la Unión de Ornitólogos del Perú (UNOP)*. 2:5. Spanish.
- Bates JM, Haffer J, Grismer E. 2004. Avian mitochondrial DNA sequence divergence across a headwater stream of the Río Tapajós, a major Amazonian river. *Journal of Ornithology*. 145:199–205.
- Bax V, Francesconi W, Quintero M. 2016. Spatial modeling of deforestation processes in the central Peruvian Amazon. *Journal for Nature Conservation*. 29:79–88.
- Begazo AJ, Valqui HT. 1998. Birds of Pacaya-Samiria National Reserve with a new population (*Myrmotherula longicauda*) and new record for Peru (*Hylophilus semicinereus*). *Bulletin of the British Ornithologists' Club*. 118:159–166.
- Bertram B. 1970. The vocal behavior of the Indian Hill mynah, *Gracula religiosa*. *Animal Behavior Monograph*. 3:79–192.
- Borges SH, Cornelius C, Ribas C, Almeida R, Guilherme E, et al. 2016. What is the avifauna of Amazonian white-sand vegetation? *Bird Conservation International*. 26:192–204.

- Capparella AP. 1987. Effects of riverine barriers on genetic differentiation of Amazonian forest undergrowth birds (Peru) [dissertation]. Baton Rouge (LA): Louisiana State University.
- Cracraft J. 1985. Historical biogeography and patterns of differentiation within the South American avifauna: Areas of endemism. In: Buckley PA, Foster MS, Morton ES, Ridgely RS, Buckley FG, editors. Neotropical ornithology. Ornithological Monographs 36. Washington (DC): American Ornithologists' Union; p. 49–84.
- Daut EF, Brightsmith DJ, Mendoza AP, Puhakka L, Peterson MJ. 2015. Illegal domestic bird trade and the role of export quotas in Peru. *Journal for Nature Conservation*. 27:44–53.
- Del-Río G, Mutchler MJ, Costa B, Hiller AE, Lima G, et al. 2021. Birds of the Jurúa River: Extensive várzea forest as a barrier to *terra firme* birds. *Journal of Ornithology*. <https://doi.org/10.1007/s10336-020-01850-0>
- Díaz-Alván J, Socolar JB, Álvarez AJ. 2017. The avifauna of the Rio Tigre basin, northern Perú. *Ornitología Neotropical*. 28:11–21.
- Dickerman RW. 1975. New subspecies of *Caprimulgus sericoaudatus* from the Amazon River Basin. *Bulletin of the British Ornithologists' Club*. 95:18–19.
- Dumont JF. 1991. Fluvial shifting in the Ucayali Depression as related to the neotectonics of the Andean foreland Brazilian Craton Border (Peru). *Géodynamique*. 6:9–20.
- Gastañaga M, Macleod R, Hennessey B, Ugarte J, Puse E, et al. 2011. A study of the parrot trade in Peru and the potential importance of internal trade for threatened species. *Bird Conservation International*. 21:76–85.
- González MOE. 1998. Birds of the lowland forest of Cerros del Sira, central Peru. *Cotinga*. 9:57–60.
- Goussard JJ. 1984. Étude préliminaire de l'avifaune du haut Ucayali et du haut Mapuya (Amazonie péruvienne) [Preliminary study of the avifauna of the upper Ucayali and upper Mapuya (Peruvian Amazon)]. *L'Oiseau et la Revue Française d'Ornithologie*. 54:305–336.
- Graham GL, Graves GR, Schulenberg TS, O'Neill JP. 1980. Seventeen bird species new to Peru from the Pampas de Heath. *Auk*. 97:366–370.
- Guilherme E, Borges SH. 2011. Ornithological records from a campina/campinarana enclave on the upper Jurúa River, Acre, Brazil. *Wilson Journal of Ornithology*. 123:24–32.
- Haffer J. 1977. Verbreitung und hybridisation der *Pionites*-papageien Amazoniens [Distribution and hybridization of the Amazonian *Pionites*-parrots]. *Bonnerzoologische Beiträge*. 28:269–278. German.
- Harvey MG, Lane DF, Hite J, Terrill RS, Figueroa S, et al. 2014a. Notes on bird species in bamboo in northern Madre de Dios, Peru, including the first Peruvian record of Acre Tody-Tyrant (*Hemitriccus cohnhaffi*). *Occasional Papers of the Museum of Natural Science, Louisiana State University*. 81:1–38.
- Harvey MG, Seeholzer GF, Cáceres AD, Winger BM, Tello JG, et al. 2014b. The avian biogeography of an Amazonian headwater: The Upper Ucayali River, Peru. *Wilson Journal of Ornithology*. 126:179–191.
- Harvey MG, Winger BM, Seeholzer GF, Cáceres AD. 2011. Avifauna of the Gran Pajonal and the southern Cerros del Sira, Peru. *Wilson Journal of Ornithology*. 123:289–315.
- Ichikawa M, Ricse A, Ugarte J, Kobayashi S. 2014. Migration patterns and land use by immigrants under a changing frontier society in the Peruvian Amazon. *Tropics*. 23:73–82.
- Johnson OJ, Howard JT, Brumfield RT. 2021. Systematics of a Neotropical clade of dead-leaf-foraging antwrens (Aves: Thamnophilidae: *Epinecrophylla*). *Molecular Phylogenetics and Evolution*. 154:106962.
- Kratter AW. 1997. Bamboo specialization by Amazonian birds. *Biotropica*. 29:100–110.
- Lane DF, Servat GP, Valqui HT, Lambert FR. 2007. A distinctive new species of tyrant flycatcher (Passeriformes: Tyrannidae: *Cnipodectes*) from southeastern Peru. *Auk*. 124:762–772.
- Lathrap DW. 1968. Occupation and changes in river channel on the central Ucayali, Peru. *American Antiquity*. 33:62–79.
- Lebbin DJ. 2007. Habitat specialization among Amazonian birds: Why are there so many *Guadua* bamboo specialists? [dissertation]. Ithaca (NY): Cornell University.
- Lebbin DJ. 2013. Nestedness and patch size of bamboo-specialist bird communities in southeastern Peru. *Condor*. 115:230–236.
- Lees AC, Moura NG, Almeida AS, Vieira ICG. 2015. Poor prospects for avian biodiversity in Amazonian oil palm. *PLOS One*. 10(5):e0122432.
- Löhr H. 1963. The use of bird calls to clarify taxonomic relationships. In: Sibley CG, editor. *Proceedings XIII International Ornithological Congress*. Baton Rouge (LA): American Ornithologists' Union; p. 544–552.
- Lowery GH Jr, O'Neill JP. 1965. A new species of *Cacicus* (Aves: Icteridae) from Peru. *Occasional Papers of the Museum of Zoology, Louisiana State University*. 33:1–5.
- Lowery GH Jr, O'Neill JP. 1966. A new genus and species of cotinga from eastern Peru. *Auk*. 83:1–9.
- Lowery GH Jr, O'Neill JP. 1969. A new species of antpitta from Peru and a revision of the subfamily Grallariinae. *Auk*. 86:1–12.
- Marra PP, Remsen JV Jr. 1997. Insights into the maintenance of high species diversity in the Neotropics: Habitat selection and foraging behavior in understory birds of tropical and temperate forests. In: Remsen JV Jr, editor. *Studies in Neotropical ornithology honoring Ted Parker*. Ornithological Monographs 48. Washington (DC): American Ornithologists' Union; p. 445–483.
- Mee A, Ohlson J, Stewart I, Wilson M, Örn P, Diaz FJ. 2002. The Cerros del Sira revisited: Birds of submontane and montane forest. *Cotinga*. 18:46–57.
- Moncrieff AE, Johnson O, Lane DF, Álvarez AJ, Balta K, et al. 2019. Avifaunal surveys along the lower Huallaga River, Region of Loreto, Peru: New distributional records, collection of topotypes, and taxonomic implications. *Wilson Journal of Ornithology*. 131:486–501.

- Mundinger PC. 1979. Call learning in the Carduelinae: Ethological and systematic considerations. *Systematic Zoology*. 28:270–283.
- Naka LN, Bechtoldt CL, Henriques LMP, Brumfield RT. 2012. The role of physical barriers in the location of avian suture zones in the Guiana Shield, northern Amazonia. *American Naturalist*. 179:E115–E132.
- Naka LN, Brumfield RT. 2018. The dual role of Amazonian rivers in the generation and maintenance of avian diversity. *Science Advances*. 4:1–13.
- Neller RJ, Salo JS, Räsänen ME. 1992. On the formation of blocked valley lakes by channel avulsion in upper Amazon foreland basins. *Zeitschrift für Geomorphologie*. 36:401–411.
- Novaes FC. 1981. A estrutura da espécie nos periquitos do gênero *Pionites* Heine (Psittacidae, Aves) [The species structure of the parrot genus *Pionites* Heine (Psittacidae, Aves)]. *Boletim do Museu Paraense Emílio Goeldi, Zoologia*. 106:1–21. Portuguese.
- O'Neill JP. 1969. Distributional notes on the birds of Peru, including twelve species previously unreported from the republic. *Occasional Papers of the Museum of Natural Science, Louisiana State University*. 37:1–11.
- O'Neill JP. 1974. The birds of Balta, a Peruvian dry tropical forest locality, with an analysis of their origins and ecological relationships [dissertation]. Baton Rouge (LA): Louisiana State University.
- O'Neill JP. 2003. Avifauna de la región de Balta, un poblado Cashinahua en el Río Curanja [Avifauna of the Balta region, a Cashinahua settlement on the Curanja River]. In: Pitman RL, Pitman N, Álvarez P, editors. *Alto Purús: Biodiversidad, Conservación y Manejo*. Lima (Peru): Impreso Gráfica and Duke University Center for Tropical Conservation; p. 97–106, 318–329. Spanish.
- O'Neill JP, Munn CA, Franke I. 1991. *Nannopsittaca dachilleae*, a new species of parrotlet from eastern Peru. *Auk*. 108:225–229.
- O'Neill JP, Pearson DL. 1974. Estudio preliminar de las aves de Yarinacocha, Departamento de Loreto, Perú [A preliminary study of the birds of Yarinacocha, Department of Loreto, Peru]. *Publicaciones del Museo de Historia Natural "Javier Prado"*, Serie A Zoología. 25:1–13. Spanish.
- Orme CDL, Davies RG, Burgess M, Eigenbrod F, Pickup N, et al. 2005. Global hotspots of species richness are not congruent with endemism or threat. *Nature*. 436:1016–1019.
- O'Shea BJ, Stotz DF, Saboya del Castillo P, Ruelas Inzunza E. 2015. Birds. In: Pitman N, Vriesendorp C, Rivera Chavez L, Wachter T, Alvira Reyes D, et al., editors. *Perú: Tapiche-Blanco. Rapid Biological Inventories Rep. 27*. Chicago (IL): Field Museum of Natural History; p. 446–471.
- Parker TA III. 1982. Observations of some unusual rainforest and marsh birds in southeastern Peru. *Wilson Bulletin*. 94:477–493.
- Parker TA III, Stotz DF, Fitzpatrick JW. 1996. Ecological and distributional databases. In: Stotz DF, Fitzpatrick JW, Moskovits DK, editors. *Neotropical birds: Ecology and conservation*. Chicago (IL): University of Chicago Press; p. 115–436.
- Parker TA III, Stotz DF, Fitzpatrick JW. 1997. Notes on avian bamboo specialists in southwestern Amazonian Brazil. In: Remsen JV Jr, editor. *Studies in Neotropical ornithology honoring Ted Parker*. *Ornithological Monographs* 48. Washington (DC): American Ornithologists' Union; p. 543–547.
- Pärssinen MH, Salo JS, Räsänen ME. 1996. River floodplain relocations and the abandonment of aborigine settlements in the upper Amazon Basin: A historical case study of San Miguel de Cunibos at the middle Ucayali River. *Geoarchaeology*. 11:345–359.
- Piana RP, Díaz AJ, Chalco LJJ, Sevillano CS, Saboya del Castillo P, Cubas PS. 2012. El Caracara Carancho (*Caracara plancus*) estaría expandiendo su rango de distribución hacia el norte y oeste de la Amazonía peruana [The Southern Caracara (*Caracara plancus*) is expanding its geographic distribution to the north and east of the Peruvian Amazon]. *Boletín de la Unión de Ornítólogos del Perú (UNOP)*. 7(2):5–11. Spanish.
- Remsen JV Jr. 1985. Community organization and ecology of birds of high elevation humid forest of the Bolivian Andes. In: Buckley PA, Foster MS, Morton ES, Ridgely RS, Buckley FG, editors. *Neotropical ornithology*. *Ornithological Monographs* 36. Washington (DC): American Ornithologists' Union; p. 733–756.
- Remsen JV Jr, Areta JL, Bonaccorso E, Claramunt S, Jaramillo A, et al. Version 9 February 2021. A classification of the bird species of South America. *American Ornithological Society*. <http://www.museum.lsu.edu/~Remsen/SACCBaseline.htm>
- Remsen JV Jr, Parker TA III. 1983. Contribution of river-created habitats to bird species richness in Amazonia. *Biotropica*. 15:223–231.
- Robinson SK. 1997. Birds of a Peruvian oxbow lake: Populations, resources, predation, and social behavior. In: Remsen JV Jr, editor. *Studies in Neotropical ornithology honoring Ted Parker*. *Ornithological Monographs* 48. Washington (DC): American Ornithologists' Union; p. 613–639.
- Rosenberg G. 1990. Habitat specialization and foraging behavior by birds of Amazonian river islands in Northeastern Peru. *Condor*. 92:427–443.
- Rowland F. 2018. eBird checklist: <https://ebird.org/checklist/S48776397>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.
- Rutt CL, Mokross K, Kaller MD, Stouffer PC. 2020. Experimental forest fragmentation alters Amazonian mixed-species flocks. *Biological Conservation*. 242:108415.
- Schmitt F, Sané R, Thibault M, Vásquez G. 2017. New locality for White-masked Antbird *Pithys castaneus* and other avian range extensions for dpto. Loreto, Peru. *Cotinga*. 39:2–11.
- Schulenberg TS, Albuja C, Rojas JI. 2006. Birds. In: Vriesendorp C, Schulenberg TS, Alverson WS, Moskovits DK, Rojas JI, editors. *Perú: Sierra del Divisor. Rapid Biological Inventories Report 17*. Chicago (IL): Field Museum of Natural History; p. 185–196.

- Schulenberg TS, Servat G. 2001. Avifauna of the northern Cordillera de Vilcabamba, Peru. In: Alonso LE, Alonso A, Schulenberg TS, Dallmeier F, editors. Biological and social assessments of the Cordillera de Vilcabamba, Peru. Washington (DC): Conservation International; p. 92–97.
- Schulenberg TS, Stotz DF, Lane DF, O'Neill JP, Parker TA III. 2010. Birds of Peru: Revised and updated edition. Princeton (NJ): Princeton University Press.
- Silva JMC, Rylands AB, Fonseca GAB. 2005. The fate of the Amazonian areas of endemism. *Conservation Biology*. 19:689–694.
- Socolar JB, Díaz-Alván J, Saboya del Castillo P, Pomara LY, O'Shea BJ, et al. 2018. Noteworthy bird records from northeastern Peru reveal connectivity and isolation in the western Amazonian avifauna. *Wilson Journal of Ornithology*. 130:94–111.
- Socolar SJ, González Ó, Forero-Medina G. 2013. Noteworthy bird records from the northern Cerros del Sira, Peru. *Cotinga*. 35:26–38.
- Srinivas A, Koh LP. 2016. Oil palm expansion drives avifaunal decline in the Pucallpa region of Peruvian Amazonia. *Global Ecology and Conservation*. 7:183–200.
- Srinivas A, Molina VJ. 2013. New distributional records of Rufous-necked Puffbird *Malacoptila rufa* and Pearly Antshrike in Peruvian Amazonia. *Cotinga*. 35:88–89.
- Sullivan BL, Wood CL, Iliff MJ, Bonney RE, Fink D, Kelling S. 2009. eBird: A citizen-based bird observation network in the biological sciences. *Biological Conservation*. 142:2282–2292.
- Terborgh J. 1985. Habitat selection in Amazonian birds. In: Cody ML, editor. Habitat selection in birds. New York (NY): Academic Press; p. 311–338.
- Terborgh J, Robinson SK, Parker TA III, Munn CA, Pierpont N. 1990. Structure and organization of an Amazonian forest bird community. *Ecological Monographs*. 60:213–238.
- Terborgh J, Weske JS. 1975. The role of competition in the distribution of Andean birds. *Ecology*. 56:562–576.
- Traylor, MA. 1958. Birds of northeastern Peru. *Fieldiana: Zoology*. 35:87–141.
- Vijay V, Reid CD, Finer M, Jenkins CN, Pimm SL. 2018. Deforestation risks posed by oil palm expansion in the Peruvian Amazon. *Environmental Research Letters*. 13:114010.
- Weir JT, Faccio MS, Pulido-Santacruz P, Barrera-Guzmán AO, Aleixo A. 2015. Hybridization in headwater regions and the role of rivers as drivers of speciation in Amazonian birds. *Evolution*. 69:1823–1834.
- Whitney BM, Alvarez AJ. 1998. A new *Herpsilochmus* antwren (Aves: Thamnophilidae) from northern Amazonian Peru and adjacent Ecuador: The role of edaphic heterogeneity of *terra firme* forest. *Auk*. 115:559–576.
- Whittaker A, Oren DC. 1999. Important ornithological records from the Rio Juruá, western Amazonia, including twelve additions to the Brazilian avifauna. *Bulletin of the British Ornithologists' Club*. 119:235–260.
- Wiley RH. 2010. Alfonso Olalla and his family: The ornithological exploration of Amazonian Peru. *Bulletin of the American Museum of Natural History*. 343:1–68.
- Williams SM. 2017. Behavioral ecology of Amazonian mixed-species flocks [dissertation]. East Lansing (MI): Michigan State University.
- Zann R. 1985. Ontogeny of the Zebra Finch distance call: I. Effects of cross-fostering to Bengalese finches. *Zeitschrift für Tierpsychologie*. 68:1–23.
- Zimmer KJ, Whittaker A, Sardelli CH, Guilherme E, Aleixo A. 2013. A new species of *Hemitriccus* tody-tyrant from the state of Acre, Brazil. In: del Hoyo J, Elliot A, Sargatal J, Christie D, editors. Handbook of the birds of the world. Special volume. New species and global index. Barcelona (Spain): Lynx Edicions; p. 292–296.

Supplemental Appendix

ML numbers in the below text refer to supporting media (photographs or audio recordings) archived online at the Macaulay Library (<https://www.macaulaylibrary.org>) and XC numbers refer to supporting audio recordings archived online at <https://www.xeno-canto.org>. LSUMZ numbers refer to catalogued specimens in the bird collection at the LSUMNS (formerly the Louisiana State University Museum of Zoology). Specimens marked with an asterisk have the round skin deposited in the bird collection of the Centro de Ornitología y Biodiversidad (CORBIDI) and are represented only by tissue samples at the LSUMNS. We follow Remsen et al. (2021) for English and scientific names of birds.

Species accounts

First records for Ucayali Region

Band-tailed Nighthawk (*Nyctiprogne leucopyga*)—We observed this species at dawn flying low over Laguna Juanacha on the mornings of 18, 21, and 22 August 2019, with ~20 individuals recorded on the 22nd. We also detected at least 3 individuals flying along a nearby stretch of the Tamaya River at dawn on 22 August 2019. We obtained audio recordings of perched individuals singing (ML176718801, ML176523901), photographs (ML176822291, ML176822261), and 2 specimens (LSUMZ 228622 and 228623*). These represent the first records for the Ucayali Region, with the nearest published record of this species from an oxbow lake in the Pacaya-Samiria National Reserve 405 km north-northeast of Laguna Juanacha (Begazo and Valqui 1998).

Individuals with distinct song types (one a deliberate “chunk-chewink”, the other a rapid, slightly ascending trill) have been found to co-occur in some localities within the Amazon Basin

(B Whitney and M Cohn-Haft, LSUMNS and Instituto Nacional de Pesquisas da Amazônia (INPA), unpubl. data), and our observations now confirm that both song types occur in Peru. Previous Peru records documented with audio recordings from the Nanay River (Álvarez and Whitney 2003; DFL, 2015: XC298511, XC298523) and the lower Javari River (B Whitney, LSUMNS, unpubl. data) all involved individuals of the “triller” song type (B Whitney, LSUMNS, pers. comm.). Our recordings were exclusively of “chunk-chewink” songs from Laguna Juanacha and represent the first documentation of this song type in Peru, although several previous records from Pacaya-Samiria without audio recordings may refer to this form. The nearest confirmed records of the “chunk-chewink” song type involve audio recordings by S Mayer and J Woude ~925 km to the east-southeast near Riberalta in Beni Department, Bolivia (XC2428 and XC437, respectively) and 8 individuals seen and heard singing and calling at multiple sites ~925 km to the northeast near the mouth of the Putumayo River in Amazonas, Brazil (M Cohn-Haft, INPA, unpubl. data). Molecular evidence demonstrated a deep divergence within currently recognized *N. leucopyga* (Sigurdsson and Cracraft 2014), but nomenclatural issues have stalled efforts to recognize 2 species. An ongoing study of the taxonomy of the genus *Nyctiprogne* seeks to unequivocally link the distinct song types to available names but requires further sampling near type localities (B Whitney and M Cohn-Haft, LSUMNS and INPA, in prep.).

White-chinned Swift (*Cypseloides cryptus*)—We detected this species at 3 headwater islands (Isla Pukani, Escondida, and Cetical), the Cerros del Sira foothills (photographed: ML137643711, ML137643761) in October 2015, and at our principal Tournavista study site in August 2019 (photographed: ML189215611), representing the first records for the Ucayali and Huánuco regions. As others have suggested (Schulenberg et al. 2010, Schmitt et al. 2017), we

believe this species is probably overlooked due to its tendency to fly high and fast overhead and the difficulty of separating it from similar species. Even distant swift flocks, however, are worth examining with patience for details in shape, flight style, and the presence of white markings on the face that can confirm this species. Voice, an underappreciated character for swift identification, is another clue.

Green-tailed Goldenthrout (*Polytmus theresiae*)—We detected 7 individuals, obtained 2 specimens (LSUMZ 228656* and 228657), and obtained an audio recording (ML176534421) at Laguna Juanacha on 22 August 2019. These records represent the first for the Ucayali Region and a range extension of ~325 km south from the nearest known site along the Blanco River (Socolar et al. 2018). In Peru, this species is known from a few patchily distributed sites across the lowlands and is generally associated with open savanna habitat (Graham et al. 1980, Schulenberg et al. 2010, Schmitt et al. 2017, Socolar et al. 2018, Moncrieff et al. 2019). At Laguna Juanacha, however, we observed this species on the edge of seasonally flooded scrub forest (~5-7 m tall) surrounding the lake and on small islands in the center of the lake.

Wire-crested Thorntail (*Discosura popelairii*)—We observed 3 individuals including a male with a long, thin crest at a flowering tree near our Kirahuanero camp 6.5 km north of Oventeni. These are the first reports of the species for the Ucayali Region and the Cerros del Sira, although the report of *Discosura* sp. at nearby Monte Tabor (Harvey et al. 2011) likely pertains to this species.

Semipalmated Plover (*Charadrius semipalmatus*)—OJ observed a single individual on 6 October 2015 at Isla Cetical with a group of Collared Plovers (*C. collaris*). The somewhat larger size and darker brown back were among features noted that distinguish this species from the much more expected Collared Plover. Semipalmated Plover is almost unknown from the

Amazonian lowlands of Peru and this report represents the first for the Ucayali Region. The nearest record east of the Andes is a photographed individual ~365 km to the southeast near Pilcopata, Madre de Dios (Zuniga 2018; ML127962681).

Killdeer (*Charadrius vociferus*)—OJ observed one individual on 6 October 2015 at Isla Cetical. The large size, dark brown upperparts, white underparts, and 2 black chest bands were plumage features noted to separate this species from other similar shorebirds. This represents the first report for the Ucayali Region and one of only a few reports for the Amazon basin. It is not clear if this individual represents an overshooting migrant of the nominate subspecies or a wanderer of the Peruvian coastal form *C. v. peruvianus*.

Wilson's Phalarope (*Phalaropus tricolor*)—We documented the first records for the Ucayali Region at Isla del Gallo on 11 October 2015 (9 individuals detected and 2 specimens collected; LSUMZ 190467 and 206900*). MGH photographed 3 individuals at Lago Yarinacocha on 20 August 2019 (ML173386201, ML173386181).

Franklin's Gull (*Leucophaeus pipixcan*)—MGH observed one adult in basic plumage at close range from a boat as it rested on driftwood floating in the Ucayali River directly in front of the port of Pucallpa on 11 December 2015. This represents the first report for the Ucayali Region and the second for the Peruvian lowlands after a November sighting of one individual in Madre de Dios (Schulenberg 1980). These reports have been followed by recent eBird records with photographs of individuals in Loreto in the months of February and March (Faherty 2017, Arévalo 2020).

Bamboo Antshrike (*Cymbilaimus sanctaemariae*)—We detected at least 4 individuals in the midstory of viny, *terra firme* forest above our first Sepahua camp (~360 m elevation) during 18-19 August 2018. We did not encounter bamboo at this locality nor did we detect other

bamboo-associated species. We obtained 2 specimens (male and female; LSUMZ 227515 and 227516*, respectively) and audio recordings from one individual (ML126272321). These represent the first records for the Ucayali Region with the nearest eBird reports from sites in the Junín (~60 km west-southwest; García-Bravo 2010) and Cuzco regions (~55 km southwest; Gamarra-Toledo 2013).

Yellow-breasted Antwren (*Herpsilochmus axillaris*)—This species was common around the town of Kirahuanero, where we collected a female (LSUMZ 206722*) on 19 September 2015 and audio-recorded one individual (ML62266401) on 21 September 2015. These represent the first records for the Ucayali Region. The song in the cited recording starts with musical introductory notes, indicative of the nominate southern subspecies and may be the northernmost locality for the taxon.

Dot-backed Antbird (*Hylophylax punctulatus*)—We collected one individual at Quebrada Caco on 9 December 2015 (LSUMZ 190525). The only previous record for this species in the Ucayali Region is a specimen collected by the Olalla brothers from the 1927 locality “Boca Rio Urubamba” (AMNH 407137), but the high catalog number (the Olalla collections from Ucayali generally have AMNH catalog numbers in the 240,000s) suggest that this bird was received by AMNH later than the regular Olalla collections and that it likely derives from the problematic collections (in terms of label accuracy) of Harvey Bassler (RH Wiley, University of North Carolina, pers. comm.; see also Wiley 2010). Thus, we consider our record from Quebrada Caco to represent the first confirmed record for the Ucayali Region.

Yellow-chinned Spinetail (*Certhiaxis cinnamomeus*)—This species was common at 2 sites in the Masisea area: at Laguna Juanacha between 18 August and 22 August 2019, where we collected 2 specimens (LSUMZ 228874* and 228875) and obtained photographs and audio

recordings (ML176232161, ML176762271, ML178720971), and the Masisea island on 23-24 August 2019, where we collected 2 specimens (LSUMZ 228876 and 228877*) and also obtained photographs and audio recordings (ML177495901, ML176831461, ML176831481). At Laguna Juanacha this species was associated with beds of aquatic *Polygonum* vegetation, whereas on the Masisea island it was found in herbaceous grassy vegetation growing out of a dried mudflat. These represent the first records for the Ucayali Region, and it was only in 2004 that the species was first conclusively documented for Peru, along the Tapiche River (Valqui and Alza 2016). It has since been reported from numerous localities in Loreto and is described as among the most common passerines along the Tapiche River (Socular et al. 2018).

Brownish Elaenia (*Elaenia pelzelni*)—We detected 7 individuals, collected one (LSUMZ 228944), and photographed 2 (ML176828481, ML176829601) at the Masisea island. These represent the first records for the Ucayali Region and a southwestward range extension of ~600 km from localities with documented individuals from the vicinity of Iquitos (Schulenberg et al. 2006).

Torrent Tyrannulet (*Serpophaga cinerea*)—GFS observed one individual at a stop on the Puerto Ocopa–Atalaya road along the Unine River ~18 km southeast of Oventeni on 9 October 2015. This represents the first report for the Ucayali Region and suggests the distribution of this species extends from the Andes (Schulenberg et al. 2010) well into the southern Cerros del Sira.

Blackish Pewee (*Contopus nigrescens*)—On 9 October 2015, GFS obtained photographs (ML137643841, ML137643821) and audio recordings (ML20446681, ML20446711) of a recently fledged Blackish Pewee at a logging track on the west side of the first major ridge of the southern Cerros del Sira travelling west from Atalaya along the Puerto Ocopa–Atalaya road (~20

km southwest of Atalaya; 880 m elevation; 10.915°S, 73.905°W). This individual was accompanied by 2 adults and a third adult was seen at another point along the track. Five days later (13 October 2015) we collected 2 specimens (LSUMZ 190600 and 206901*) at this same locality and obtained additional audio recordings (ML2219302, ML60912731). In central Peru, Blackish Pewee is known from the northern Cerros del Sira (Socolar et al. 2013) and from photographs by DFL from Abra Aguachine in the Pasco Region (ML68654731, ML68654741). Our records from 2015 are the first for the Ucayali Region and presently the southernmost locality for Andean populations.

The striking vocal similarity between Blackish Pewee and nominate Tropical Pewee (*Contopus cinereus cinereus*) strongly suggests that the Blackish/Tropical Pewee complex merits detailed taxonomic reassessment. There are 3 main vocal groups within Tropical Pewee: 1) the “Northern/*bogotensis* group” of Middle and northern South America, 2) the “Tumbes/*punensis* group” of western Ecuador and Peru, and 3) the “Southern/*cinereus* group” of central South America, generally south and east of Amazonia (see Farnsworth et al. 2020). The “Tumbes/*punensis* group” has already been recognized at the species level by some authorities based on its distinctive vocalizations (Gill et al. 2020, Ridgely and Greenfield 2001). We suspect that all 3 groups likely involve species-level taxa and that Blackish Pewee may be closely allied to, if not best considered conspecific with, the Southern/*cinereus* group of Tropical Pewee. Furthermore, it may be of interest to compare the Brazilian Shield and Andean populations presently considered Blackish Pewee to see if they are in fact sister populations. At present there is no named taxon to exclusively represent the disjunct Brazilian shield population, and the current situation of it belonging to the same taxon as the Peruvian Andean population (*C. n. canescens*) seems biogeographically unlikely.

Amazonian Black-Tyrant (*Knipolegus poecilocercus*)—We collected one female (LSUMZ 228952) in low scrub next to a small, swampy oxbow near Laguna Juanacha on 21 August 2019. This represents the first record for the Ucayali Region. The nearest record appears to be a female specimen collected by the Olalla brothers in 1927 near Sarayacu, Loreto Department, approximately 230 km to the north-northwest (Wiley 2010).

Gray-chested Greenlet (*Hylophilus semicinereus*)—This species was fairly common at Laguna Juanacha, where we audio-recorded (ML176759331), photographed (ML176815081), and collected 3 individuals (LSUMZ 228977, 228978*, 228979*) between 18 and 22 August 2019 in seasonally flooded scrub forest surrounding the laguna. AEM also heard one individual in taller flooded forest with a scrubby understory at Cocha Larga on 22 August 2019. These are the first records for the Ucayali Region and represent a range extension of ~320 km south from the nearest documented population along the Blanco River in the Loreto Region (O’Shea et al. 2015).

Southern Martin (*Progne elegans*)—We observed up to ~10 individuals of this species on 24-25 July 2019 flying high over the Hotel Ecológico Manish in Pucallpa and DFL obtained photographs (ML169594871, ML169594881). These represent the first records for the Ucayali Region of this likely overlooked austral migrant.

First records for Huánuco Region

White-chinned Swift (*Cypseloides cryptus*)—See account in *First records for Ucayali Region* section above.

Needle-billed Hermit (*Phaethornis philippii*)—AEM observed one individual on 29 August 2019 in the interior of a forest fragment at our principal Tournavista study site. Although

there are specimen records at sites nearby around Pucallpa (Traylor 1958; American Museum of Natural History (AMNH) series from 59 km west of Pucallpa; LSUMNS series from Yarinacocha), this represents the first report for the Huánuco Region.

Variegated Flycatcher (*Empidonomus varius*)—We photographed one individual in second-growth/edge habitat at our principal Tournavista field site on 28 August 2019 (ML176834521). This represents the first record for the Huánuco Region. We detected another individual near our Preferida camp in the Ucayali Region ~70 km east-northeast on 19 August 2019. In Peru this species is known as an austral migrant and winter resident (Schulenberg et al. 2010), but it occurs in Bolivia as an austral migrant and breeder (Chesser 1997, Herzog et al. 2016). There is now some evidence, however, that suggests the species may also breed locally in foothills of southern Peru; there are several documented records around San Pedro, at ~1400 m elevation, on the Kosñipata Road in the Cuzco Region from October through January (DFL pers. obs., Drucker 2013; ML77662851, ML77662771, ML128669661, ML238994251, ML239003631). The species is largely found in Amazonian lowlands in the austral winter but breeds at similar elevations (up to 1700 m, DFL pers. obs.) as near as La Paz department, Bolivia, only ~400 km to the southeast.

First records for Junín Region

Black-and-white Hawk-Eagle (*Spizaetus melanoleucus*)—GFS photographed an adult 6 km northeast of Puerto Ocopa (1100 m elevation; ML137644151). This widespread species is uncommon throughout its range, and this record represents the first for the Junín Region, although this species is previously reported from the adjoining Cerros del Sira in the Ucayali Region (Harvey et al. 2011).

Undulated Antshrike (*Frederickena unduliger*)—AEM observed and audio-recorded (ML127733621) a male on 28 September 2018 in *terra firme* forest ~9.5 km east of Mayapo. This represents the first record for the Junín Region. The nearest previous record is of 2 specimens (AMNH 239047 and 239048) ~70 km north-northwest at Lagarto in the Ucayali Region (Zimmer 1944).

Manu Antbird (*Cercomacra manu*)—We detected populations of this species in a *Guadua* bamboo tract ~30 km northwest of Sepahua and at Nuevo San Martín and Mayapo; we collected 2 specimens at each locality (LSUMZ 227541* and 227542*, 227539 and 227540, 227543 and 227544*, respectively) and obtained several audio recordings (ML126986681, ML127230241). Our Mayapo records represent the first for the Junín Region, and in combination with our Ucayali records, help fill in a gap between its distribution in the southeastern lowlands of Peru (Schulenberg et al. 2006) and an outlying record from ~50 km northwest of Atalaya (Harvey et al. 2014b).

Brown-rumped Foliage-gleaner (*Automolus melanopezus*)—We detected this bamboo-associated species at 4 different localities, collected 4 specimens across 3 localities (LSUMZ 227592, 227593*, 227594*, 227595), and obtained audio recordings (ML127291971, ML127833331). A sound-vouchered specimen (LSUMZ 227595, ML127833331) from Mayapo represents the first record for the Junín Region. Previous records in the Ucayali Region were limited to specimens from Balta (O'Neill 2003), so our records establish a more widespread presence of the species.

Striolated Manakin (*Machaeropterus striolatus*)—We audio-recorded one individual (ML127821961) and collected 3 (LSUMZ 227694*, 227696, 227697) in foothill forest above the town of Cushireni. We collected an additional individual across the Tambo River at Mayapo

(LSUMZ 227695*). These records, the southernmost for the species (see Guilherme 2013), represent the first for the Junín Region and a range extension of ~90 km south from records along the Cohengua River (Harvey et al. 2014b).

Black-backed Tody-Flycatcher (*Poecilatriccus pulchellus*)—We collected 2 males (LSUMZ 227721* and 227722) in dense, viny, second-growth habitat next to agricultural fields at Mayapo. These represent the first records in the Junín Region for this Peruvian endemic and a range extension of ~100 km northwest from the nearest known locality near Camisea, Cuzco Region (Schulenberg et al. 2006).

Shiny Cowbird (*Molothrus bonariensis*)—We observed and photographed an adult male at a small wetland 10 km south of Atalaya on the Puerto Ocopa–Atalaya road on 8 October 2015 (ML234524811). Surprisingly, this appears to constitute the first record for the Junín Region of this widespread species of open and disturbed habitats throughout South America.

Other notable distributional records

Black-bellied Whistling-Duck (*Dendrocygna autumnalis*)—Schulenberg et al. (2010) described the status in Peru as “rare and local in [Peruvian] Amazonia, where mostly found in the north.” We found this species on several occasions on the middle Ucayali just upriver of Pucallpa, and returning to Pucallpa by boat on the evening of 24 August 2019 we counted 65 individuals at one location (photographed: ML176832781, ML176832771). This, in addition to other records from the Pucallpa area, including one of 30 individuals (Angulo 2017; ML68592231), the Cohengua River and Atalaya (the first records for the Ucayali Region; Harvey et al. 2014b), San Juan de Inuya (Herrera-Peralta 2016; ML29894141), and other sites in

southern Ucayali and Madre de Dios (Cuadros 2016), suggests that the species may occur regularly in the central Peruvian Amazon.

Brazilian Teal (*Amazonetta brasiliensis*)—We observed 3 individuals on Isla del Gallo near Atalaya on 11 October 2015 and photographed this species (ML170303251, ML176838181) at Yarinacocha on 4 visits during July to September 2019, with up to ~12 individuals present on 31 August 2019. Reports of this species in Peru were until recently restricted to the Madre de Dios Region, where the species is probably increasing (Schulenberg et al. 2010). Based on our records, along with recent photographed eBird records in Ucayali (Carrasco and Osorio 2019, Custodio 2019) and the first documented records for the San Martín Region involving 3 individuals in 2019 (DFL photographs: ML165990011, ML165990021), we suspect that occurrences are increasing in the central Peruvian lowlands and farther north. The first Ucayali records (a pair) of this species were only documented in May 2011 by MGH on a river island near Atalaya (Harvey et al. 2014b).

Silky-tailed Nightjar (*Antrostomus sericocaudatus*)—We detected this poorly known species regularly in the interior of *terra firme*-like forest at our Preferida camp. We obtained audio recordings (ML176677051, ML176688391), collected one female on 19 August 2019 (LSUMZ 228626), and found one active nest. The nest was a scrape on the ground containing small leaf parts, twigs, and bark peel, and was situated on the edge of a treefall gap (15 by 8 m) with no canopy cover. Although numerous nests have been documented in Madre de Dios to the south (Wilkinson 2009), this is the first nest documented in the Ucayali Region. The nearest records of the species are a specimen from 1946 collected at Yarinacocha (FMNH 185571; Traylor 1958) and the type specimen of *A. s. mengeli* from 1972 collected 59 km east of Pucallpa (Dickerman 1975).

Lesser Violetear (*Colibri cyanotus*)—MGH observed one individual visiting a *terra firme* forest stream near our Preferida camp on 6 August 2019. The combination of violet auriculars, green throat, and blue-green tail distinguished it from similar lowland hummingbird species. This is typically an Andean species, with the nearest records from the Cerros del Sira (Mee et al. 2002, Harvey et al. 2011), and its presence in the lowlands may have been tied to a cold front (*friaje*) affecting the area at the time.

Ocellated Crane (*Micropygia schomburgkii*)—We heard one individual in a *pajonal* 3 km north of Oventeni on 20 September 2015, and audio-recorded one individual in a *pajonal* along the Puerto Ocopa–Atalaya road on 13 October 2015 at 1060 m elevation (10.894°S, 74.104°W; ML22195761). This latter locality is ~3 km west-northwest of the only previous report of this species in the Ucayali Region by H van Oosten, who recorded multiple individuals (XC152544) on 28 September 2013. Together, these reports suggest the regular presence of this species in the Gran Pajonal. The nearest records are ~235 km to the southeast in foothill grasslands of the Cuzco Region at ~1250-1500 m elevation (Robbins et al. 2011). Farther abroad, populations are known from the Mayo Valley (~600 km to the northwest; first recorded near Moyobamba by B Whitney in August 2013) and the Pampas del Heath (~605 km to the southeast; Graham et al. 1980).

American Golden-Plover (*Pluvialis dominica*)—We photographed one individual (ML189218931, ML176837991) on 31 August and 1 September 2019 at Lago Yarinacocha. The only previous record in the Ucayali Region is from Balta (O’Neill 2003).

Buff-breasted Sandpiper (*Calidris subruficollis*)—We observed and photographed one individual at Lago Yarinacocha on 20 August 2019 (ML173386131) and again on 31 August 2019 (ML176836491, ML176836481). Although regular in the Iquitos area, this species is

poorly known elsewhere in the Peruvian Amazon. Previous specimen records in the Ucayali Region are from Balta (16 September 1964; LSUMZ 33976) and 59 km west of Pucallpa (30 October 1971; AMNH 822202).

Puna Ibis (*Plegadis ridgwayi*)—AEH photographed up to 4 individuals together (ML228040501, ML228040511) along the Urubamba River in the mudflats next to our camp ~30 km northwest of Sepahua. This is the second record of this vagrant to the Ucayali Region after a sighting of 4 flyovers near the Cohengua River (Harvey et al. 2014b). Observers of any *Plegadis* in the Peruvian Amazon should try to obtain the best documentation possible, because it is conceivable that White-faced Ibis (*P. chihi*) might also rarely occur in the area.

White-tailed Hawk/Variable Hawk (*Geranoaetus albicaudatus/polyosoma*)—We observed 2 individuals of *Geranoaetus* in the Gran Pajonal. The first was a subadult flying over the road between Oventeni and Kirahuanero on 3 July 2015 (photographed: ML284610121, ML284610131, ML284610141). Several features of this bird point to Variable Hawk, including the whitish throat and long wings relative to their width. This potentially represents the first record of Variable Hawk for the Ucayali Region, but due to limitations of the photographs not all authors feel confident in identifying this individual to species. A second individual with notably more advanced plumage was observed by GFS on 3 October 2015 in a pasture 3 km north of Oventeni (10.710°S, 74.233°W) with connectivity to larger *pajonales*. This individual had a white throat and underparts with a grey head (giving it a hooded appearance), reddish wing coverts, and a mottled white tail with a dark terminal band. The mottling on the tail and upperparts suggest it was in advanced subadult plumage. Although these plumage features combined point towards White-tailed Hawk, we lack conclusive documentation and only later,

after extensive review of the photographs of the first individual, noted the possibility of Variable Hawk in the same area.

The geographic distributions of these species in Peru differ substantially, but they both are typically found in open habitats. Variable Hawk is widespread on the Peruvian coast and up to 4600 m elevation in the Andes, but only down to 2000 m elevation on the east slope (Schulenberg et al. 2010). White-tailed Hawk was first observed in Peru in the Pampas del Heath (Graham et al. 1980), but this species appears to be fairly widespread, occurring in various arid intermontane valleys (Robbins et al. 2011, Valqui and Alza 2016) and in areas where lowland forest is being cleared for pasture such as along the Interoceanic Highway near Puerto Maldonado, Madre de Dios (e.g., Alferez 2016; ML28310501). The nearest observations to the Gran Pajonal were made by GFS and MGH at Satipo in the Junín Region on multiple occasions from 2008-2011.

Brown-banded Puffbird (*Notharchus ordii*)—We photographed (ML177762021, ML177762001, ML177761981) one individual in tall, *terra firme*-like forest at Preferida on 12 August 2019. This is the second locality for this species in the Ucayali Region, following records from the Cohengua River ~200 km to the south (Harvey et al. 2014b). Although often associated with white-sand forests (Schulenberg. et al. 2010, Álvarez et al. 2013), our observation along with those in Loreto (Socolar et al. 2018) indicate this species can be found in a variety of poor-soil habitats.

Spotted Puffbird (*Bucco tamatia*)—We photographed (ML176808301, ML177757161) and collected 2 specimens (LSUMZ 228676* and 228677) of this species at Preferida, where it was fairly common. This represents the second locality for the Ucayali Region after a previous record from the Cohengua River (Harvey et al. 2014b). Schulenberg et al. (2010) depicted a

range gap between records in Loreto and Madre de Dios; with the species now known from multiple localities in Ucayali, we suggest this gap was likely the result of insufficient sampling.

Rufous-necked Puffbird (*Malacoptila rufa*)—This species was common at our Masisea camp, where we collected 6 individuals (LSUMZ 228680, 228681, 228682*, 228683*, 228684, and 228685*) and obtained photographs (ML176806471) and audio recordings (ML176520621). This represents only the second known locality in the Ucayali Region and the first east of the Ucayali River. The previous records involve photographed individuals from a banding project in a forest fragment ~20 km southwest of Pucallpa (Srinivas and Molina 2013; ML188749141, ML188747881).

Blue-cheeked Jacamar (*Galbula cyanicollis*)—We collected one female (LSUMZ 227500) in *terra firme* forest ~30 km northwest of Sepahua on 6 September 2018. This extends the known range for this species in Peru ~80 km southeast of records from the Cohengua River (Harvey et al. 2014b).

Crested Caracara (*Caracara plancus*)—We photographed single individuals on the lower Urubamba River on 16 August 2018 (ML126205131, ML126205121, ML126205111), in Masisea on 20 August 2019 (ML176817011), and near Tournavista on 28 August 2019 (ML176834931). Additionally, we detected 2 individuals at Breu on 14-17 December 2015. These records further support an assessment of rapid range expansion for this species as deforestation progresses in the Amazonian lowlands of Peru (Piana et al. 2012, Socolar et al. 2018).

Stripe-chested Antwren (*Myrmotherula longicauda*)—We audio-recorded a single individual at Cocha Larga on 22 August 2019 (ML176778461) in viny, seasonally flooded forest. This species is generally known from higher elevations, in foothill forest between 500 and

1550 m elevation (Schulenberg et al. 2010), but apparently disjunct lowland populations have been documented in similar blackwater habitats in Loreto: at Pacaya-Samiria National Reserve by Begazo and Valqui (1998), and near San José de Vinuya and the Tapiche River (in association with bamboo scrub) by Socolar et al. (2018). There are also lowland records of this species from northeastern Madre de Dios (Harvey et al. 2014a) in second-growth *terra firme* forest near extensive *Guadua* bamboo. Although the songs we recorded at Cocha Larga sounded like typical *M. longicauda*, due to the lack of specimen material from this and the other lowland sites listed above, we cannot rule out the possibility that a new taxon may be involved. This species should be a target of future collecting efforts in the area.

Short-billed Leaf-tosser (*Sclerurus ruficularis*)—We collected 3 individuals at our Masisea camp (LSUMZ 228830*, 228831, 228832*). The only other known locality for this species in the Ucayali Region is along the Cohengua River (Harvey et al. 2014b).

Black-faced Cotinga (*Conioptilon mcilhennyi*)—We detected 3 individuals and collected one male (LSUMZ 227705) in the canopy of hilly *terra firme* forest on the north bank of the Inuya River near Nuevo San Martín on 20 September 2018. AEM also audio-recorded 2 individuals (ML126274851) in hilly *terra firme* forest near our first Sepahua camp on 18 August 2018. Although this species has been associated with floodplain forest (Ridgely and Tudor 1994, Schulenberg et al. 2010), our observations along with those in Acre, Brazil (Whittaker and Oren 1999) and Madre de Dios (Harvey et al. 2014a) indicate that this species also regularly occurs in *terra firme* forests.

Cinereous Becard (*Pachyramphus rufus*)—We found two birds during 7–9 December 2015 at Quebrada Caco and two more on 23 August 2019 at the Masisea island where we photographed (ML176829251) and collected 2 specimens (LSUMZ 228903* and 228904). This

species was previously known in the Ucayali Region from a 1962 specimen (LSUMZ 28454) collected by JP O'Neill at Yarinacocha and a few later eBird reports by others at the same locality (e.g., Angulo and Valqui 2017; ML68602021). Based on this group of records, it appears that this species is uncommon but regular in scrubby habitats well up the middle Ucayali River.

Rufous Twistwing (*Cnipodectes superrufus*)—We audio-recorded (ML127846901, ML127852231) and collected a pair (LSUMZ 227723* and 227724) at Mayapo on 1 October 2018. Although thought to be a bamboo obligate (Lane et al. 2007, Tobias et al. 2008), we observed this pair in young second-growth forest with viny tangles near a stream. We note, however, there were tracts of bamboo in the vicinity. Rufous Twistwing is a species of conservation concern (Tobias et al. 2008), listed by the IUCN as vulnerable (BirdLife International 2017), but its presence in a matrix of cleared, cultivated land suggests that this species may tolerate moderate levels of disturbance—indeed, its preferred habitats (bamboo) require a certain level of disturbance. Previous reports from the Junín Region of this poorly known species include sight records ~15 km to the southeast (García-Bravo 2010) and an audio-recording by D García-Olaechea (XC44574) ~35 km to the southeast.

Foothill Elaenia (*Myiopagis olallai*)—We detected 2 individuals and obtained photographs (ML137644111) and audio recordings (ML20448231) of one along the Puerto Ocopa–Atalaya road (780 m elevation; 10.899°S, 74.195°W) on 9 October 2015 and obtained a single specimen (LSUMZ 190564) on 13 October 2015. In 2012, DFL and others detected this species at the same site, and, in nearby Junín, obtained photographs (ML225110261) and recordings (ML225131591).

Foothill Elaenia was first described from a small series of specimens from foothill forest on the east slope of the Andes near Zamora, Ecuador (Coopmans and Krabbe 2000). Two

additional subspecies were described from higher elevation humid Andean forest in Antioquia, Colombia and Perijá, Venezuela (Cuervo et al. 2014). In their description of the species, Coopmans and Krabbe (2000) included a single Peruvian specimen (AMNH 819936) among those they identified as Foothill Elaenia. This specimen was from the Cordillera Vilcabamba (Cuzco Region) and had been previously identified as the similar Forest Elaenia (*M. gaimardii*). The Peruvian records of Foothill Elaenia likely pertain to the nominate subspecies, based on the slower pace of the long song and overall yellow, not grey, underparts. The specimen we collected is the second of Foothill Elaenia from Peru and the first with associated tissue samples. Over 800 km separate records in Junín, Ucayali, and Cuzco from the next nearest records in Zamora-Chinchipe Province, Ecuador. Extensive coverage of appropriate elevations between these records suggest that, rather than an artifact of sampling, there may indeed be a large gap in this species' distribution.

Sooty-headed/Yungas Tyrannulet (*Phyllomyias griseiceps/weedeni*)—AEM audio-recorded one individual (ML126263771) in an overgrown *chacra* near our first Sepahua camp on 17 August 2018. This species complex is essentially unknown in the lowlands of Peru, with the only additional records from around Iberia in Madre de Dios (Harvey et al. 2014a). There are also audio recordings by D Buzzetti in nearby Acre, Brazil (XC490282, XC490600, XC490605). Other records in central Peru involved foothill birds in the Gran Pajonal (Harvey et al. 2011) and Cuzco (Robbins et al. 2011). Harvey et al. (2014a) noted a cline in vocalizations between populations of *P. griseiceps* in central and southeast Peru and populations of *P. weedeni* in southernmost Peru and Bolivia, which complicates identification. Based on vocal differences, we suspect that nominate *P. griseiceps* is probably best considered a separate species from cis-Andean populations in Ecuador and Peru, which are likely more closely related to *P. weedeni*.

Further taxonomic work is needed to sort out the relationships within this species complex, which also includes the Planalto Tyrannulet (*Phyllomyias fasciatus*).

Southern Scrub-Flycatcher (*Sublegatus modestus*)—We photographed multiple individuals (ML177773811, ML177773851, ML176812101, ML176812111) and collected one (LSUMZ 228949) at Preferida between 12 August and 20 August 2019. We also photographed individuals at the Masisea island on 23-24 August 2019 (ML176828501, ML176829801). This species is described as a “rare austral migrant” in southern Amazonia by Schulenberg et al. (2010), but our records suggest that it is a regular migrant that has perhaps simply been overlooked.

Yellow-browed Tyrant (*Satrapa icterophrys*)—We collected 6 individuals in a recently burned *pajonal* north of Oventeni on 20 September 2015 (LSUMZ 190601, 190602, 190603, 206741*, 206742*, and 206744*). In Peru, this species is an austral migrant known principally from riverine habitats in Madre de Dios with scattered reports from open habitats in the foothills and a specimen from 2200 m elevation near Sandia, Puno on 4 November 1980 collected by L Binford (LSUMZ 98498). The first record of this species for Ucayali was from a river island opposite Atalaya (Harvey et al. 2014b). Our records constitute the first specimens for the Ucayali Region and are the most northerly records for this species in Peru.

Pied Water-Tyrant (*Fluvicola pica*)—We collected the first 2 specimens for Peru at Lago Yarinacocha on 19 December 2015 (LSUMZ 190606 and 206937*), and an additional specimen at the Masisea island on 24 August 2019 (LSUMZ 228953). In July to August 2019 we also photographed individuals at multiple wetlands along the middle Ucayali River (ML189211691, ML176823931, ML170303971) and at a roadside wetland in the Tournavista area (ML176833161). This species was only recently accepted as a vagrant on the Peruvian bird

list based on photographs from December 2009 at Yarinacocha and Contamana (Loreto Region; Angulo et al. 2015). The presence of records in both the austral winter and summer, along with photographs of an individual in juvenal plumage at Yarinacocha (ML170237761) and pair building a nest (Carrasco and Walker 2018; ML101708551), suggests that this species is a breeding resident in the central lowlands of Peru rather than an irregular migrant from northern South America, as initially speculated (Angulo et al. 2015). The paucity of records for this species between central Peru and central Colombia (first Ecuador record in 2007 and still very few overall; Freile et al. 2013), suggests that the central Peru population is truly disjunct. Although eBird records with photographs of this species have increased exponentially in the Pucallpa area in the last few years (e.g., Delzo 2017, Angulo 2019), it is unclear whether this is due to a recent range expansion or increased coverage and awareness among ornithologists and birdwatchers.

A complicating factor in assessing the status of Pied Water-Tyrant in Peru is that reports did not generally include, until recently, documentation of the distinguishing features separating this species from the Black-backed Water-Tyrant (*Fluvicola albiventer*). For decades, many authorities considered Black-backed Water-Tyrant a subspecies of Pied Water-Tyrant (Hellmayr 1927, Zimmer 1937, Meyer de Schauensee 1970, AOU 1998, Dickinson 2003), and it was not until the 1990s that these taxa began to receive treatment as separate species (Sibley and Monroe 1990, Ridgely and Tudor 1994). Furthermore, it was assumed, until recently, that Peruvian records of *Fluvicola* all belonged to Black-backed Water-Tyrant (Schulenberg et al. 2010), given that it was the only form of the complex documented by specimens prior to these recent reports (T Schulenberg, Cornell Lab of Ornithology, pers. comm.). Black-backed Water-Tyrant is documented from the eastern lowlands of Peru during the austral winter based on an adult female

specimen (NHMUK 1888.1.1.142) collected at Pebas, Loreto Region, on 4 June 1866 (photographs of specimen examined; Sclater 1888, Sclater and Salvin 1867) and an individual photographed in the Madre de Dios Region on 25 August 2020 (ML257962491; Maceda 2020).

Given that sight records of *Fluvicola* in Amazonian Peru prior to the split of Black-backed and Pied water-tyrants did not provide details allowing identification to species, it remains difficult to assess the status of the 2 species in the area. O’Neill and Pearson (1974) mentioned that “*Fluvicola pica*” was observed regularly in small numbers around the edge of Lago Yarinacocha, but it is unclear to which taxon they were referring. Nevertheless, we suspect that O’Neill and Pearson (1974) observed Pied Water-Tyrant, and that the central Peru population of this species has been present for a substantial period of time. We hope that, henceforth, reports of *Fluvicola* in Amazonian Peru will be documented well enough to provide a better understanding of the status of both species in the country.

Citron-bellied Attila (*Attila citriniventris*)—We obtained 2 specimens (LSUMZ 228961* and 228962) and an audio recording (ML176673191) in *terra firme* forest near the Preferida camp, and heard individuals on most days at this site. This is the second locality for this species in the Ucayali Region, following records from the Cohengua River (Harvey et al. 2014b). This species is often associated with white-sand forests (Schulenberg et al. 2010, Álvarez et al. 2013), which are not present at this locality. However, the *terra firme* forest where we detected this species was on relatively poorly drained soil with swampy terrain, which apparently also meets the habitat requirements of this species (Álvarez et al. 2013).

Scaly-breasted Wren (*Microcerculus marginatus*)—We detected this species at multiple localities on both sides of the Ucayali River and in the Tambo–Urubamba interfluvium. Our observations are consistent with previous findings of 2 distinctive song types in western

Amazonia: one found north of the Amazon and west of the Ucayali and another found south of the Amazon and east of the Ucayali (Ridgely and Tudor 1989, Schulenberg et al. 2010, Harvey et al. 2014b). As expected, we recorded the northern song type at Tournavista (ML176795751) and the southern song type at Masisea (ML176675911). We found that south of Atalaya the 2 song types appear to replace each other across the Tambo River, with the northern song type recorded at Cushireni (left bank; ML240625471) and an individual seen by AEM in the vicinity of and shortly after hearing a southern song type at Serjali (right bank). AEM also recorded an apparent southern song type of Scaly-breasted Wren at Mayapo (right bank; ML240620741). However, due to the potential for confusion between the southern song type of Scaly-breasted Wren and the song of Wing-banded Wren (*M. bambla*) and our lack of visuals of singing Scaly-breasted Wren in this area, the song type in the Tambo–Urubamba interfluvium remains to be confirmed.

Genetic data indicate that the song types in Amazonia belong to distinct clades that diverged ~4 Ma (Smith et al. 2014). The name *M. m. marginatus*, described from “Bogotá” (Sclater 1855), is associated with the northern song type, leaving no currently recognized subspecies to assign to the southern song type. We recommend resurrecting the name *M. m. bolivianus* (Carriker 1935), which had been synonymized with *M. m. marginatus* by Peters (1960), to recognize the southern song type as a distinct taxon. The complex southern song type is the only one known throughout the range of the species in Bolivia, so it can be safely associated with the name *M. m. bolivianus*.

Bicolored Conebill (*Conirostrum bicolor*)—We detected a total of 3 individuals, photographed one (ML176828461, ML176828471), and collected another (LSUMZ 229014) at the Masisea island on 23 August 2019. The only previous record in the Ucayali Region is a

specimen collected by J Schunke from “Pucallpa” on 4 February 1956 (University of Arizona Bird Collection 017889). Presumably this individual was collected from an island in the Ucayali River given the species’ close association with river islands (Rosenberg 1990) and its absence from surveys of the Yarinacocha area (O’Neill and Pearson 1974).

Literature cited

- Alferez J. 2016. eBird checklist: <https://ebird.org/checklist/S29472677>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.
- Álvarez A J, Metz MR, Fine PVA. 2013. Habitat specialization by birds in western Amazonian white-sand forests. *Biotropica*. 45:365–372.
- Álvarez A J, Whitney BM. 2003. New distributional records of birds from white-sand forests of the northern Peruvian Amazon, with implications for biogeography of northern South America. *The Condor*. 105:552–566.
- Angulo F. 2017. eBird checklist: <https://ebird.org/peru/checklist/S38965692>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.
- Angulo F, Valqui H T. 2017. eBird checklist: <https://ebird.org/checklist/S39128958>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.
- Arévalo G. 2020. eBird checklist: <https://ebird.org/checklist/S64352845>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.
- Begazo AJ, Valqui H T. 1998. Birds of Pacaya-Samiria National Reserve with a new population (*Myrmotherula longicauda*) and new record for Peru (*Hylophilus semicinereus*). *Bulletin of the British Ornithologists’ Club*. 118:159–166.

- Carrasco Z G, Osorio D. 2019. eBird checklist: <https://ebird.org/checklist/S61488483>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.
- Carrasco Z G, Walker B. 2018. eBird checklist: <https://ebird.org/checklist/S45969690>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.
- Carriker Jr MA. 1935. Descriptions of new birds from Bolivia, with notes on other little-known species. *Proceedings of the Academy of Natural Sciences of Philadelphia*. 87:313–341.
- Chesser RT. 1997. Patterns of seasonal and geographical distribution of austral migrant flycatchers (Tyrannidae) in Bolivia. In: Remsen JV Jr, editor. *Studies in Neotropical ornithology honoring Ted Parker*. *Ornithological Monographs* 48. Washington D.C.: American Ornithologists' Union; p. 171–204.
- Cuadros S. 2016. Registro del Pato-Silbón de Vientre Negro (*Dendrocygna autumnalis*) en la estación biológica Cocha Cashu y revisión de su distribución en Perú [Record of Black-bellied Whistling-Duck (*Dendrocygna autumnalis*) at the Cocha Cashu biological station and a revision of its distribution in Peru]. *Boletín de la Unión de Ornitólogos del Perú (UNOP)*. 11(1):73–79. Spanish.
- Cuervo AM, Stiles FG, Lentino M, Brumfield RT, Derryberry EP. 2014. Geographic variation and phylogenetic relationships of *Myiopagis olallai* (Aves: Passeriformes; Tyrannidae), with the description of two new taxa from the northern Andes. *Zootaxa*. 3873:1–24.
- Custodio O. 2019. eBird checklist: <https://ebird.org/checklist/S62819374>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.

- Dickerman RW. 1975. New subspecies of *Caprimulgus sericocaudatus* from the Amazon River Basin. *Bulletin of the British Ornithologists' Club*. 95:18–19.
- Dickinson EC. 2003. *The Howard and Moore complete checklist of the birds of the world*. 3rd ed. London (UK): Christopher Helm. 1040 p.
- Drucker J. 2013. eBird checklist: <https://ebird.org/checklist/S16382631>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.
- Faherty M. 2017. eBird checklist: <https://ebird.org/checklist/S36001839>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.
- Farnsworth A, del Hoyo J, Lebbin DJ, Collar N, Kirwan GM. 2020. Tropical Pewee (*Contopus cinereus*), version 1.0. In: Billerman SM, Keeney BK, Rodewald PG, Schulenberg TS, editors. *Birds of the world*. Ithaca (NY): Cornell Lab of Ornithology. DOI: <https://doi.org/10.2173/bow.tropew1.01>
- Gamarra-Toledo V. 2013. eBird checklist: <https://ebird.org/checklist/S19791518>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.
- García-Bravo A. 2010. eBird checklist: <https://ebird.org/peru/checklist/S16292795>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.
- Gill F, Donsker D, Rasmussen P. 2020. *IOC World Bird List (v10.1)* [cited 13 May 2020]. DOI: <https://doi.org/10.14344/IOC.ML.10.1>.
- Graham GL, Graves GR, Schulenberg TS, O'Neill JP. 1980. Seventeen bird species new to Peru from the Pampas de Heath. *The Auk*. 97:366–370.
- Guilherme E. 2013. A range extension of Striped Manakin *Machaeropterus striolatus*

- (Bonaparte, 1838) (Aves: Pipridae) based on records from southwestern Brazilian Amazon. *Check List*. 9(6):1530–1532.
- Harvey MG, Lane DF, Hite J, Terrill RS, Figueroa S, et al. 2014a. Notes on bird species in bamboo in northern Madre de Dios, Peru, including the first Peruvian record of Acre Tody-Tyrant (*Hemitriccus cohnhafti*). *Occasional Papers of the Museum of Natural Science, Louisiana State University*. 81:1–38.
- Harvey MG, Seeholzer GF, Cáceres A D, Winger BM, Tello JG, et al. 2014b. The avian biogeography of an Amazonian headwater: The Upper Ucayali River, Peru. *Wilson Journal of Ornithology*. 126:179–191.
- Harvey MG, Winger BM, Seeholzer GF, Cáceres A D. 2011. Avifauna of the Gran Pajonal and the southern Cerros del Sira, Peru. *Wilson Journal of Ornithology*. 123:289–315.
- Herrera-Peralta K. 2016. eBird checklist: <https://ebird.org/checklist/S30104114>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.
- Herzog SK, Terrill RS, Jahn AE, Remsen JV Jr, Maillard O, et al. 2016. *Birds of Bolivia: field guide*. Santa Cruz de la Sierra (Bolivia): Asociación Armonía.
- Maceda K. 2020. eBird checklist: <https://ebird.org/peru/checklist/S72820874>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.
- Mee A, Ohlson J, Stewart I, Wilson M, Örn P, Diaz F J. 2002. The Cerros del Sira revisited: birds of submontane and montane forest. *Cotinga*. 18:46–57.
- Moncrieff AE, Johnson O, Lane DF, Álvarez A J, Balta K, et al. 2019. Avifaunal surveys along the lower Huallaga River, Region of Loreto, Peru: new distributional records, collection of topotypes, and taxonomic implications. *Wilson Journal of Ornithology*. 131:486–501.

- O'Neill JP. 2003. Avifauna de la región de Balta, un poblado Cashinahua en el Río Curanja [Avifauna of the Balta region, a Cashinahua settlement on the Curanja River]. In: Pitman RL, Pitman N, Álvarez P, editors. Alto Purús: Biodiversidad, Conservación y Manejo. Lima (Peru): Impreso Gráfica and Duke University Center for Tropical Conservation; p. 97–106, 318–329. Spanish.
- O'Neill JP, Pearson DL. 1974. Estudio preliminar de las aves de Yarinacocha, Departamento de Loreto, Perú [A preliminary study of the birds of Yarinacocha, Department of Loreto, Peru]. Publicaciones del Museo de Historia Natural "Javier Prado", Serie A Zoología: 1–13. Spanish.
- O'Shea BJ, Stotz DF, Saboya del Castillo P, Ruelas Inzunza E. 2015. Birds. In: Pitman N, Vriesendorp C, Rivera Chavez L, Wachter T, Alvira Reyes D, et al., editors. Perú: Tapiche-Blanco. Rapid Biological Inventories Rep. 27. Chicago (IL): Field Museum of Natural History; p. 446–471.
- Peters JL. 1960. Check-list of birds of the world. Mayr E, Greenway Jr JC, editors. Cambridge (MA): Museum of Comparative Zoology. 506 p.
- Piana R P, Díaz A J, Chalco L JJ, Sevillano CS, Saboya del Castillo P, Cubas P S. 2012. El Caracara Carancho (*Caracara plancus*) estaría expandiendo su rango de distribución hacia el norte y oeste de la Amazonía peruana [The Southern Caracara (*Caracara plancus*) is expanding its geographic distribution to the north and east of the Peruvian Amazon]. Boletín de la Unión de Ornitólogos del Perú (UNOP). 7(2):5–11. Spanish.
- Remsen JV Jr, Areta JI, Bonaccorso E, Claramunt S, Jaramillo A, et al. Version 9 February 2021. A classification of the bird species of South America. American Ornithological Society. <http://www.museum.lsu.edu/~Remsen/SACCBaseline.htm>

- Ridgely RS, Greenfield PJ. 2001. The Birds of Ecuador. Vol. 1. Status, distribution, and taxonomy. Ithaca (NY): Cornell University Press.
- Ridgely RS, Tudor G. 1989. The birds of South America, volume 1: The oscine passerines. Austin (TX): University of Texas Press.
- Ridgely RS, Tudor G. 1994. The birds of South America, volume 2: The suboscine passerines. Austin (TX): University of Texas Press.
- Robbins MB, Geale D, Walker B, Davis TJ, Combe M, et al. 2011. Foothill avifauna of the upper Urubamba Valley, dpto. Cusco, Peru. *Cotinga*. 33:41–52.
- Rosenberg G. 1990. Habitat specialization and foraging behavior by birds of Amazonian river islands in Northeastern Peru. *The Condor*. 92:427–443.
- Schmitt F, Sané R, Thibault M, Vásquez G. 2017. New locality for White-masked Antbird *Pithys castaneus* and other avian range extensions for dpto. Loreto, Peru. *Cotinga*. 39:2–11.
- Schulenberg TS. 1980. A Franklin's Gull (*Larus pipixcan*) in southeastern Peru. *Gerfaut*. 70:403–404.
- Schulenberg TS, Stotz DF, Rico L. 2006. Distribution maps of the birds of Peru. Version 1.0. Environment, Culture & Conservation (ECCo), The Field Museum [cited 1 May 2020]. http://fm2.fieldmuseum.org/uw_test/birdsofperu
- Schulenberg TS, Stotz DF, Lane DF, O'Neill JP, Parker TA III. 2010. Birds of Peru: revised and updated edition. Princeton (NJ): Princeton University Press.
- Sclater PL. 1855. On the birds received in the collections from Santa Fé di [sic] Bogotá. *Proceedings of the Zoological Society of London*. p. 131–164.

- Sigurdsson S, Cracraft J. 2014. Deciphering the diversity and history of New World nightjars (Aves: Caprimulgidae) using molecular phylogenetics. *Zoological Journal of the Linnean Society*. 170:506–545.
- Smith BT, McCormack JE, Cuervo AM, Hickerson MJ, Aleixo A, et al. 2014. The drivers of tropical speciation. *Nature*. 515:406-409.
- Socolar JB, Díaz-Alvan J, Saboya del Castillo P, Pomara LY, O’Shea BJ, et al. 2018. Noteworthy bird records from northeastern Peru reveal connectivity and isolation in the western Amazonian avifauna. *Wilson Journal of Ornithology*. 130:94–111.
- Socolar SJ, Gonzalez O, Forero-Medina G. 2013. Noteworthy bird records from the northern Cerros del Sira, Peru. *Cotinga*. 35:26–38.
- Srinivas A, Molina V J. 2013. New distributional records of Rufous-necked Puffbird *Malacoptila rufa* and Pearly Antshrike in Peruvian Amazonia. *Cotinga*. 35:88–89.
- Traylor, MA. 1958. Birds of northeastern Peru. *Fieldiana: Zoology*. 35:87–141.
- Valqui T, Alza L. 2016. Analisis de especımenes de cuatro especies de aves nuevas para el Peru depositados en la coleccion cientıfica de CORBIDI [Analysis of specimens of four species of birds new for Peru deposited in the scientific collection of CORBIDI]. *Boletın de la Union de Ornitologos del Peru*. 11(2):29–41. Spanish.
- Wiley RH. 2010. Alfonso Olalla and his family: the ornithological exploration of Amazonian Peru. *Bulletin of the American Museum of Natural History*. 343:1–68.
- Wilkinson FA. 2009. Observations on the breeding biology of the Silky-tailed Nightjar (*Caprimulgus sericocaudatus mengeli*). *Wilson Journal of Ornithology*. 121:498–505.
- Zimmer JT. 1944. Studies of Peruvian birds. No. 49. Notes on *Frederickena* and *Ochthoeca*. *American Museum Novitates*. 1263:1–5.

Zuniga H. 2018. eBird checklist: <https://ebird.org/peru/checklist/S50591236>. eBird: An online database of bird distribution and abundance. Ithaca (NY): Cornell Lab of Ornithology.